

iMedix: Your Personal Health Advisor.

Cialis Sublingual

When Cialis Sublingual starts working, the effect will be proceeding for more than a day. Usually, it goes on for 24 – 36 hours. Other medicaments work only for 8-10 hours. The difference is quite obvious.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Different erectile disorders is one of the most common problems in the whole range of male diseases. In Massachusetts there was conducted a Male Aging Study, which proves that about 40% of men in the age of 40 have encountered erectile dysfunction. It spreads with time and by the age of 70 there are more than 70% of men facing this disease. In such cases, it's either about hoping for the best or always having a backup plan. This backup plan is also called Tadalafil, the inhibitor, used for the treatment of erectile dysfunction. But you cannot buy it its original form, it's only available in pills like Cialis or Viagra. Comparing these two medicines, it's safe to say, that first one has higher efficiency, takes less time to start working and gives more lasting effect. Tadalafil relaxes vessels in the penis and increases the amount of blood in it, causing the erection. Although, it does not raise the libido, so first make sure to wake up your sexual desire. At the end of 2011 US Food and Drug Administration (FDA) approved Cialis. It was declared a cure for symptoms of erectile dysfunction and benign prostatic hyperplasia. Cialis usually goes in small yellow pills of 2.5 mg, 5 mg, 10 mg and 20 mg in the almond shape. First, you put it under the tongue and watch it quickly dissolving. After that it goes straight to the bloodstream, skipping the digestive system. Due to this, it takes only 20-30 minutes for the tablet to start working (some even claim 16 minutes). Other medicines in this category need about 60 minutes to kick in. Cialis is a prescribed drug, which means that you can only start taking it after the conversation with your doctor. In this way, it's more likely to avoid side effects. A doctor can also explain to you what dosage you should take, what time period is more suitable, what kind of food or drink can interact with the pill. Some people would try taking more of it, but it wouldn't help to increase its efficiency. Vice versa, it can harm your health and cause some serious problems. It's better to take Sublingual Cialis on an empty stomach to ensure its performance. If you're going for bigger dose, like 20 mg, use it before sex. In case you want the stable effect, try lowering the dose to 2.5 mg or 5 mg, so you can take it on the daily basis. As for precautionary measures, it is reasonable not to take this drug, while having grapefruit juice, alcohol or tobacco. Cialis, accompanied by tobacco may cause sudden optic-nerve related vision loss, called NAION. So if you do smoke, make sure to tell it to your doctor, before using this medication. Another thing, you have to warn him about, is an abnormality of genitals. Headaches, diarrhea, nausea and flu symptoms are the most common side effects of Generic Cialis Sublingual. If you have any of this, don't panic, it will probably go on its own. Not that common effects are blurred vision, low blood pressure, and painful or prolonged erection. If you experience any of these symptoms, immediately call for medical help. This way you will avoid possible damage to your organism. Taking bigger dosage, than prescribed one increases the chances of having these

symptoms and will even make it more severe. Study's has also shown that index of side effects may increase in men after 65 years.

Side Effects

Uses

Interactions

- -
 -
 -
-

Other Details
