

# iMedix: Your Personal Health Advisor.

## Desonide Ointment

---

Desonide ointment is a low-potency corticosteroid used to treat skin conditions such as eczema, psoriasis, and dermatitis. It reduces inflammation and immune responses, alleviating redness, itching, and swelling. The ointment form boosts skin hydration and is particularly suited for sensitive areas like the face, minimizing the risk of side effects like skin thinning.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

## Description

Side Effects Stinging, burning, itching, irritation, dryness, or redness may occur when first applied to the skin. This should disappear in a few days as your body adjusts to the medication. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: stretch marks, skin thinning/discoloration, acne, excessive hair growth, small red bumps on the skin (folliculitis). Skin infections can become worse when using this medication. Tell your doctor promptly if redness, swelling or irritation does not improve. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat a variety of skin conditions (such as eczema, dermatitis, allergies, rash). Desonide reduces the swelling, itching and redness that can occur in these types of conditions. Desonide is a mild corticosteroid. How to use desonide topical Use this medication only on the skin. Wash your hands. Before applying the medication, clean and dry the affected area. Apply a thin film of the medication to the affected area and gently rub in as directed by your doctor, usually 2 to 3 times daily. If using the gel, apply the gel usually only 2 times daily. Do not bandage, wrap, or cover the area unless you are instructed to do so by your doctor. If used in the diaper area on an infant, do not use tight-fitting diapers or plastic pants. After applying the medication, wash and dry your hands, unless you are using this medication to treat the hands. When applying this medication near the eyes, avoid getting it in the eyes because this may worsen or cause glaucoma, or cause irritation. Also, avoid getting this medication in the nose or mouth. If you get this medication in your eyes, nose, or mouth, rinse right away with plenty of water. Use this medication only for the condition prescribed. Tell your doctor if your condition lasts or gets worse after 2 weeks. Precautions Before using desonide, tell your doctor or pharmacist if you are allergic to it; or to other corticosteroids (such

as hydrocortisone, prednisone); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: poor blood circulation, diabetes, immune system problems, other skin conditions (such as rosacea, perioral dermatitis). Do not use if there is an infection or sore in the area to be treated. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Children may be more sensitive to the effects of too much corticosteroid medication. Consult your doctor for more details. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication, when applied to the skin, passes into breast milk. However, similar drugs, when taken by mouth, pass into breast milk. Consult your doctor before breast-feeding. If used during breastfeeding, avoid applying the medication to the breast/nipple area to prevent the infant from swallowing the medication. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. This medication may be harmful if swallowed. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for other skin problems unless told to do so by your doctor. A different medication may be necessary in those cases. Tell all your doctors you use or have used this medication. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Different brands/strengths of this medication may have different storage requirements. Read the package labeling or ask your pharmacist for the storage requirements for the product you are using. Store at room temperature. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---