

# iMedix: Your Personal Health Advisor.

## Hydroquinone-Vit C-Vit E Ac Cream

---

Search for medical details concerning hydroquinone-vitamin C-vitamin E acetate topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Mild burning, stinging, redness, and dryness may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has prescribed this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Stop using hydroquinone and tell your doctor right away if you have any serious side effects, including: blistering, skin cracking, blue-black darkening of the skin. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Hydroquinone is used to lighten the dark patches of skin (also called hyperpigmentation, melasma, “liver spots,” “age spots,” freckles) caused by pregnancy, birth control pills, hormone medicine, or injury to the skin. This medicine works by blocking the process in the skin that leads to discoloration. How to use Hydroquinone-Vit C-Vit E Ac Cream Follow all directions on the product package, or use as directed by your doctor. Before using, apply a small amount of this medicine to an area of unbroken skin, and check the area within 24 hours for any serious side effects. If the test area is itching, red, puffy, or blistering, do not use this product and contact your doctor. If there is just mild redness, then treatment with this product may begin. Apply this medication to the affected areas of skin, usually twice daily or as directed by your doctor. This medication is for use on the skin only. If it is used incorrectly, unwanted skin lightening may occur. Avoid getting this product in your eyes or on the inside of your nose or mouth. If you do get this medication in those areas, flush with plenty of water. This medication may make the treated areas of skin more sensitive to the sun. Avoid prolonged sun exposure, tanning booths, and sunlamps. Use a sunscreen and wear protective clothing on the treated areas of skin when outdoors. Use this medication regularly to get the most benefit from it. To help you remember, use it at the same times each day. Tell your doctor if your condition lasts or gets worse after 2 months. Precautions Before using hydroquinone, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients (such as sulfites), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or

pharmacist your medical history, especially of: asthma, other skin conditions (such as eczema, psoriasis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms after swallowing may include: shaking (tremors), seizures. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another skin problem unless told to do so by your doctor. A different medication may be necessary in that case. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

