

iMedix: Your Personal Health Advisor.

Rapivab Vial

Search for medical details concerning Rapivab (PF) intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects This drug usually has no side effects. If you have any unusual effects, contact your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. The flu itself or peramivir may rarely cause serious mental/mood changes. This may be more likely in children. Tell your doctor right away of any signs of unusual behavior, including: confusion, hallucinations, agitation, self-injury. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** Peramivir is used to treat symptoms caused by the flu virus (influenza) if you have had symptoms for 2 days or less. It helps make the symptoms (such as stuffy nose, cough, sore throat, fever/chills, aches, tiredness) less severe and shortens the recovery time by about 1 to 2 days. This medication works by stopping the flu virus from growing. It is not a substitute for the flu vaccine. (**How to use Rapivab Vial** This medication is given as a single dose as directed by your doctor. It is given slowly into a vein, usually over 15 to 30 minutes. The dosage is based on your medical condition, age, and response to treatment. Children's dosage is also based on weight. Tell your doctor if your condition lasts or gets worse, or if new symptoms appear. **Precautions** Before using peramivir, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease (including dialysis treatment). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. **Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a

provincial poison control center. Do not share this medication with others. This medication is not a substitute for the flu vaccine. Talk with your doctor about the risks and important benefits of getting a yearly flu shot. Not applicable. Not applicable. This medication is given in a hospital or clinic and will not be stored at home. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Tell your doctor if you have received flu vaccine in the nose within 2 weeks before treatment with this medication. This medication may lower your protection from flu vaccine given in the nose. Wait at least 2 days after ending treatment with this medication before receiving flu vaccine given in the nose.

Side Effects

Uses

Interactions

Other Details
