

# iMedix: Your Personal Health Advisor.

## VISINE ALLERGY RELIEF Drops

---

Search for medical details concerning Visine-AC ophthalmic (eye) on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Stinging/redness in the eye, widened pupils, or blurred vision may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: eye pain, worsening redness/itching/swelling in or around the eyes, other vision problems, shakiness (tremor), fast/pounding/ irregular heartbeat, headache, sweating, weakness, nervousness. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How long does it take for this medicine to take effect? The onset of action for VISINE ALLERGY RELIEF Drops is relatively rapid. The effects can be noticed within minutes of application. Pheniramine maleate, as an antihistamine, and naphazoline hydrochloride, as a vasoconstrictor, begin to work soon after topical application to the eye, providing symptomatic relief. How long do the effects of this medicine last? The duration of the effects of VISINE ALLERGY RELIEF Drops can vary but generally lasts for several hours. Users may experience relief from allergy symptoms for 4 to 6 hours, necessitating reapplication for continued symptom control throughout the day, depending on individual response and the severity of symptoms. Is it safe to consume alcohol while taking this medicine? There is no direct interaction between the topical application of VISINE ALLERGY RELIEF Drops and the consumption of alcohol. However, alcohol consumption does not impact the efficacy or safety profile of the eye drops when used as directed. Is this a habit forming medicine? VISINE ALLERGY RELIEF Drops are not known to be habit-forming. The active ingredients do not possess addictive properties and are used for symptomatic relief of eye allergies rather than inducing psychological or physiological dependency. Can this medicine be taken during pregnancy? The safety of VISINE ALLERGY RELIEF Drops during pregnancy has not been established. While topical ophthalmic application results in minimal systemic absorption, caution is advised. Pregnant women should consult with a healthcare provider before using these eye drops to ensure that the benefits outweigh any potential risks. Can this medicine be taken while breast-feeding? Given

the minimal systemic absorption of the active ingredients in VISINE ALLERGY RELIEF Drops through topical ophthalmic application, the risk to nursing infants is likely low. However, as with pregnancy, nursing mothers should consult with a healthcare provider to evaluate the safety of using these eye drops during lactation. Uses Tetrahydrozoline is a decongestant used to relieve redness in the eyes caused by minor eye irritations (such as smog, swimming, dust, or smoke). It belongs to a class of drugs known as sympathomimetic amines. It works by temporarily narrowing the blood vessels in the eye. Some brands of tetrahydrozoline eye drops may contain lubricants. Lubricants help protect the eyes from more irritation and dryness. How to use VISINE ALLERGY RELIEF Drops Follow all directions on the product package, or use as directed by your doctor. If you have any questions, ask your doctor or pharmacist. To apply eye drops, wash your hands first. To avoid contamination, do not touch the dropper tip or let it touch your eye or any other surface. Remove contact lenses before applying the eye drops. Wait at least 10 minutes after using this medication before inserting contact lenses. Before using, check this product visually. Do not use if the liquid has changed color or is cloudy. Discard the medication away from children and pets. Use this medication in the affected eye(s) up to 4 times a day as needed or as directed by your doctor. Tilt your head back, look upward, and pull down the lower eyelid to make a pouch. Hold the dropper directly over the eye and place 1 drop into the pouch. Look downward and gently close your eyes for 1 to 2 minutes. Place one finger at the corner of the eye near the nose and apply gentle pressure. This will prevent the medication from draining away from the eye. Try not to blink and do not rub the eye. Repeat these steps if your dose is for more than 1 drop and for the other eye if so directed. Do not rinse the dropper. Replace the dropper cap tightly after each use. If you are using another kind of eye medication (such as drops or ointments), wait at least 5 minutes before applying other medications. Use eye drops before eye ointments to allow the eye drops to enter the eye. Overuse of this type of medication may result in increased eye redness (rebound effect when drug wears off). If this occurs, tell your doctor. Do not use this medication for longer than 3 to 4 days at a time. If your condition lasts or gets worse after 72 hours, or if you think you may have a serious medical problem, stop using this medication and get medical help right away. Precautions Before using tetrahydrozoline, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients (such as preservatives like benzalkonium chloride), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: glaucoma, heart problems (such as heart attack, chest pain), high blood pressure, diabetes, eye infection/injury, overactive thyroid (hyperthyroidism). After you apply this drug, your vision may become temporarily blurred. Do not drive, use machinery, or do any activity that requires clear vision until you can do it safely. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include drowsiness, severe sweating, decreased body temperature, slow/shallow breathing. If your symptoms are due to allergies, avoid allergens that may be the cause of your symptoms. Some common causes of allergies include pollen, grasses, weeds, dust mites, and pet dander. If your doctor has directed you to use this medication on a regular schedule and you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---