

iMedix: Your Personal Health Advisor.

Aciphex

Search for medical details concerning AcipHex oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Headache may occur. If this effect lasts or gets worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: symptoms of a low magnesium blood level (such as muscle spasms, irregular heartbeat, seizures), signs of lupus (such as rash on nose and cheeks, new or worsening joint pain). This medication may rarely cause a severe intestinal condition due to a bacteria called *C. difficile*. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. Rarely, proton pump inhibitors (such as rabeprazole) have caused vitamin B-12 deficiency. The risk is increased if they are taken every day for a long time (3 years or longer). Tell your doctor right away if you develop symptoms of vitamin B-12 deficiency (such as unusual weakness, sore tongue, or numbness/tingling of the hands/feet). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing, signs of kidney problems (such as change in the amount of urine). This is not a complete list of possible side effects. If you notice other effects, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** Rabeprazole is used to treat certain stomach and esophagus problems (such as acid reflux, ulcers). It works by decreasing the amount of acid your stomach makes. It relieves symptoms such as heartburn, difficulty swallowing, and cough. This medication helps heal acid damage to the stomach and esophagus, helps prevent ulcers, and may help prevent cancer of the esophagus. Rabeprazole belongs to a class of drugs known as proton pump inhibitors (PPIs). **How to use Aciphex** Read the Medication Guide and the Patient Information Leaflet if available from your pharmacist before you start taking rabeprazole and each time you get a refill. If you have any questions, ask your doctor or pharmacist. If you are using the tablets, take your dose by mouth with or without food as directed by your doctor, usually 1 to 2 times daily. Swallow the tablet whole with water. Do not crush, chew, or split the tablet. Doing so can release all of the drug at once, increasing the risk of side effects. If you are

using the capsules, take the dose 30 minutes before a meal as directed by your doctor, usually once daily. Do not swallow the capsule whole. Open the capsule and sprinkle the contents onto a small amount of soft food (such as applesauce or yogurt) or liquid. The food or liquid that you use should be at or below room temperature. Swallow the entire mixture within 15 minutes of preparing it. Do not chew or crush the prepared mixture. The dosage and length of treatment are based on your medical condition and response to treatment. In children, the dosage is also based on weight. If needed, antacids may be taken along with this medication. If you are also taking sucralfate, take rabeprazole at least 30 minutes before sucralfate. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time(s) each day. Continue to take this medication for the prescribed length of treatment even if you are feeling better. Tell your doctor if your condition lasts or gets worse. The risk of side effects goes up over time. Ask your doctor how long you should take this medication. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. If your doctor instructs you to use this medication regularly for a long time, lab and/or medical tests (such as a magnesium blood test, vitamin B-12 levels) may be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. A product that may interact with this drug is: methotrexate (especially high-dose treatment). Some products need stomach acid so that the body can absorb them properly. Rabeprazole decreases stomach acid, so it may change how well these products work. Some affected products include ampicillin, atazanavir, erlotinib, levoketoconazole, nelfinavir, pazopanib, rilpivirine, sparsentan, certain azole antifungals (itraconazole, ketoconazole, posaconazole), among others. This medication may interfere with certain lab tests, possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug. Does Aciphex interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Precautions Before taking rabeprazole, tell your doctor or pharmacist if you are allergic to it; or to similar drugs (such as lansoprazole, omeprazole); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, lupus. Some symptoms may actually be signs of a more serious condition. Get medical help right away if you have: heartburn with lightheadedness/sweating/dizziness, chest/jaw/arm/shoulder pain (especially with shortness of breath, unusual sweating), unexplained weight loss. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Proton pump inhibitors (such as rabeprazole) may increase your risk for bone fractures, especially with longer use, higher doses, and in older adults. Talk with your doctor or pharmacist about ways to prevent bone loss/fracture, such as by taking calcium (such as calcium citrate) and vitamin D supplements. Older adults may be more sensitive to the side effects of this drug, especially bone loss and fractures (see above), and *C. difficile* infection (see Side Effects section). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. However, similar drugs pass into breast milk. The effects on a nursing infant are unknown. Consult your doctor before breast-feeding.

Side Effects

Uses

Interactions

Other Details
