

iMedix: Your Personal Health Advisor.

Acne Cleansing Bar

Search for medical details concerning Acne Cleansing Bar topical on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Skin reactions such as peeling, itching, irritation, and reddened skin may occur, especially at the start of treatment. If any of these effects last or get worse, tell your doctor or pharmacist promptly. You may need to apply smaller amounts of the drug or use it less often. Consult your doctor or pharmacist for more details. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Benzoyl peroxide products are usually used to treat mild to moderate acne. When used to treat acne, benzoyl peroxide works by reducing the amount of acne-causing bacteria and by causing the skin to dry and peel. Some brands are used to treat a certain skin condition called rosacea. Benzoyl peroxide may help to decrease symptoms caused by rosacea, such as pimples and bumps on the face. Check the ingredients on the label even if you have used the product before. The manufacturer may have changed the ingredients. Also, products with similar names may contain different ingredients meant for different purposes. Taking the wrong product could harm you. **How to use Acne Cleansing Bar** Read the Patient Information Leaflet if available from your pharmacist before you start using this product and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Before using this product to treat acne for the first time, check if you are allergic to it. To check, put a small amount on one or two small areas of acne for three days. If you have an allergic reaction, stop using this product and get medical help right away (see the symptoms of an allergic reaction in the Side Effects section). If you don't have an allergic reaction, you can use the product. There are many benzoyl peroxide products available. Many can be purchased without a prescription. Some products (such as creams, cleansers, foam, or lotions) may require a prescription. Consult your doctor or pharmacist on the choice of the product that is best for you. If you are using the over-the-counter product to self-treat, read and follow all directions on the product package before using this medication. If you have any questions, consult your pharmacist. If your doctor has prescribed this medication, use it as directed. Some products require shaking before use.

Check your product package to see if your form of this medication needs to be shaken. If you are using this medication to treat acne, apply benzoyl peroxide to the areas of your skin affected by acne. If you are using this medication to treat rosacea, apply benzoyl peroxide to your face. Avoid getting this medication into your eyes, inside your nose or mouth, on the lips, or on any areas of broken skin since it may cause irritation. If this occurs, flush the area with plenty of water. Wash your hands after applying this product. Benzoyl peroxide may bleach hair or fabrics. Use carefully, and avoid contact with hair, clothing, and furnishings. If using cleansers containing benzoyl peroxide, wet the affected area. Gently rub the cleanser into the skin for 10-20 seconds. Work into a full lather and rinse thoroughly and then pat dry. If too much drying occurs, you may need to rinse the cleanser off sooner or use it less often. Consult your doctor or pharmacist for further details. If you are using the liquid wash, cleansing pad, or cleanser bar, use it instead of soap once or twice a day. Wet the skin before applying, gently work up a lather for 10 to 20 seconds, and rinse thoroughly. Do not scrub the skin. If using the shaving cream, first wet the area to be shaved. Apply a small amount of the medication, gently rub into the area, and then shave. If you are using a cream, foam, lotion, or gel, apply it to the affected area as directed, usually once or twice a day. Before applying each dose, gently wash the affected area with a mild cleanser, then pat dry. Apply a small amount of medicine to the affected area and rub in gently. Some products may require rinsing off after application for a certain amount of time. Check your product package to see if your form of this medication needs to be rinsed off. If you are using the medicated pads or sponges, wash the affected skin with a mild cleanser, then pat dry. Wipe the pad or sponge gently onto the skin to apply the medication, usually once or twice a day to the skin as directed. The dosage is based on your medical condition and response to therapy. Improvement in acne or rosacea is usually seen after 3 to 4 weeks of use, and maximum benefit after 8 to 12 weeks of use. Do not increase your dose or use this drug more often or for longer than directed. Your condition will not improve any faster, and your risk of side effects will increase. If your condition lasts or gets worse, or if you think you may have a serious medical problem, get medical help right away. Precautions Before using benzoyl peroxide, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients (such as perfumes), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication may make you more sensitive to the sun. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned or have skin blisters/redness. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Keep all regular medical and lab appointments. If you are using this product on a regular schedule and miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. A product that may interact with this drug is: dapsone used on the skin.

Side Effects

Uses

Interactions

Other Details
