

iMedix: Your Personal Health Advisor.

Acuvail Dropperette, Single Use Dropper

Search for medical details concerning Acuvail (PF) ophthalmic (eye) on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects This medication may sting or burn your eyes for 1-2 minutes when applied. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: eye pain/swelling, vision changes (such as decreased vision), bleeding inside the eye. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

Uses This medication is used to treat eye pain and swelling (inflammation) after cataract surgery. Ketorolac belongs to a class of drugs known as nonsteroidal anti-inflammatory drugs (NSAIDs). It works by blocking certain natural substances in your body to reduce pain and swelling.

How to use Acuvail Dropperette, Single Use Dropper Apply this medication to the affected eye as directed by your doctor, usually twice daily starting 1 day before surgery, continuing on the day of surgery, and then for 2 weeks after surgery. To apply eye medication, wash your hands first. To avoid contamination, do not touch the dropper tip or let it touch your eye or any other surface. Do not wear contact lenses while using this medication. Sterilize contact lenses according to the manufacturer's directions, and check with your doctor before you begin using them again. Tilt your head back, look upward, and pull down the lower eyelid to make a pouch. Hold the dropper directly over your eye and place one drop into the pouch. Look downward, gently close your eyes, and place one finger at the corner of your eye (near the nose). Apply gentle pressure for 1 to 2 minutes before opening your eyes. This will prevent the medication from draining out. Try not to blink or rub your eye. Wait several minutes for your vision to clear before driving or operating machinery. For single-use vials, open the vial right before use. After use, discard any remaining solution right away. Do not keep an opened vial for later use. If you are using another kind of eye medication (such as drops or ointments), wait at least 5 minutes between applying each medication. Use eye drops before eye ointments to allow the drops to enter the eye. Do not use this medication more often or for longer than directed by your doctor. This medication is usually used only for a short time. Prolonged use may slow or

delay healing and may increase your chance of serious side effects. Tell your doctor if your condition lasts or if it gets worse (for example, you develop worsening eye pain/itching/swelling). Precautions Before using ketorolac, tell your doctor or pharmacist if you are allergic to it; or to aspirin; or to other NSAIDs (such as ibuprofen, celecoxib); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: asthma (including a history of worsening breathing after taking aspirin or other NSAIDs), bleeding problems, recent multiple eye surgeries, other eye problems (such as dry eye syndrome, cornea problems), diabetes, rheumatoid arthritis, nasal polyps. After you apply this drug, your vision may become temporarily blurred or unstable. Do not drive, use machinery, or do any activity that requires clear vision until you can do it safely. If you develop an eye infection or injury, ask your doctor if you should continue to use this medication. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk, but it is unlikely to harm a nursing infant. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another eye condition unless told to do so by your doctor. A different medication may be necessary in that case. Lab and/or medical tests (such as eye exams) should be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Store the single-use vials in the pouch provided and fold the pouch ends closed. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug are: corticosteroids (such as prednisone). This medication may increase the risk of bleeding when used with other drugs that also may cause bleeding. Examples include anti-platelet drugs such as clopidogrel, "blood thinners" such as dabigatran/enoxaparin/warfarin, among others. Check all prescription and nonprescription medicine labels carefully since many medications contain pain relievers/fever reducers (aspirin, NSAIDs such as ibuprofen or naproxen). These drugs are similar to ketorolac and may increase your risk of side effects if taken together. However, if your doctor has directed you to take low-dose aspirin to prevent heart attack or stroke (usually 81-162 milligrams a day), you should continue taking the aspirin unless your doctor instructs you otherwise. Ask your doctor or pharmacist for more details.

Side Effects

Uses

Interactions

Other Details
