

iMedix: Your Personal Health Advisor.

Albenza

Search for medical details concerning Albenza oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Nausea, vomiting, abdominal pain, headache, or temporary hair loss may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: severe headache that doesn't go away, vision changes, yellowing eyes/skin, severe stomach/abdominal pain, dark urine, unusual tiredness, easy bruising/bleeding, signs of infection (such as sore throat that doesn't go away, fever, chills), signs of kidney problems (such as change in the amount of urine), confusion, very stiff neck. Get medical help right away if you have any very serious side effects, including: seizures. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This medication is used to treat certain tapeworm infections (such as neurocysticercosis and hydatid disease). **How to use Albenza** Take this medication by mouth with meals as directed by your doctor, usually 1 to 2 times daily. If you or your child have trouble swallowing tablets, you may crush or chew your dose and take it with water. For some conditions (such as hydatid disease), your doctor may direct you to take this medication in a treatment cycle (for example, twice daily with meals for 28 days then stopping the medication for 2 weeks). Carefully follow your doctor's instructions. The dosage is based on your weight, medical condition, and response to treatment. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time(s) each day. If your doctor directs you to take this medication in a treatment cycle, it may be helpful to mark the calendar with a reminder of when you need to restart the medication. Continue to take this medication until the full prescribed amount is finished, even if symptoms disappear after a few days. Stopping the medication too early may result in a return of the infection. Other medications (such as corticosteroids, anti-seizure medications) may also be prescribed for your current condition. Take those medications exactly as directed by your doctor. Avoid eating grapefruit or drinking grapefruit juice while using this medication unless your doctor or pharmacist says you may do so safely. Grapefruit can increase the

chance of side effects with this medicine. Ask your doctor or pharmacist for more details. Tell your doctor if your condition lasts or gets worse. Precautions Before taking albendazole, tell your doctor or pharmacist if you are allergic to it; or to other benzimidazole anthelmintic drugs (such as mebendazole); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, biliary tract problems (such as blockage), blood/bone marrow disorders. This medication may cause liver problems. Because drinking alcohol increases the risk of liver problems, limit alcoholic beverages while using this medication. Check with your doctor or pharmacist for more information. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant or plan to become pregnant. You should not become pregnant while using albendazole. Albendazole may harm an unborn baby. Your doctor should order a pregnancy test before you start this medication. Ask about reliable forms of birth control while using this medication and for 3 days after the last dose. If you become pregnant, talk to your doctor right away about the risks and benefits of this medication. This medication passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to. Lab and/or medical tests (such as liver function tests, complete blood counts) should be done before you start taking this medication and while you are taking it. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions See also How to Use section. Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
