

iMedix: Your Personal Health Advisor.

Allergy Eye (Naphazoline-Pheniramine) 0.025 %-0.3 % Drops

Search for medical details concerning Allergy Eye (naphazoline-pheniramine) ophthalmic on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Stinging/redness in the eye, widened pupils, or blurred vision may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: eye pain, worsening redness/itching/swelling in or around the eyes, vision problems, shakiness (tremor), fast/pounding/ irregular heartbeat, headache, sweating, weakness, nervousness. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to temporarily relieve eye redness, puffiness, itching, and watering that commonly occur with allergies. Naphazoline is a decongestant. It belongs to a class of drugs known as sympathomimetic amines. It works by temporarily narrowing the blood vessels in the eye. Pheniramine is an antihistamine. It works by blocking certain natural substances (histamines) that cause allergic symptoms. Do not use this medication to treat redness and irritation from wearing contact lenses. How to use Allergy Eye (Naphazoline-Pheniramine) 0.025 %-0.3 % Drops To apply eye drops, wash your hands first. To avoid contamination, do not touch the dropper tip or let it touch your eye or any other surface. Remove contact lenses before applying the eye drops. Wait at least 10 minutes after using this medication before inserting contact lenses. If your eyes are red, ask your doctor whether you should wear your contact lenses. Before using, check this product visually. Do not use if the liquid has changed color or is cloudy. Discard the medication away from children and pets. Use this medication in the affected eye(s) up to 4 times a day as needed or as directed by your doctor. Tilt your head back, look upward, and pull down the lower eyelid to make a pouch. Hold the dropper directly over the eye and place 1 drop into the pouch. Look downward and gently close your eyes for 1 to 2 minutes. Place one finger at the corner of the eye near the nose and apply gentle pressure. This will prevent the medication from draining away from the eye. Try not to blink and do not rub the eye. Repeat these steps if your dose is for more than 1 drop and for the other eye if so directed. Do not rinse the dropper. Replace the dropper cap tightly after each use. If

you are using another kind of eye medication (such as drops or ointments), wait at least 5 minutes before applying other medications. Use eye drops before eye ointments to allow the drops to enter the eye. Overuse of this type of medication may result in increased eye redness (rebound effect when drug wears off). If this occurs, tell your doctor. Do not use this medication for longer than 3 to 4 days at a time. If your condition lasts or gets worse after 72 hours, or if you think you may have a serious medical problem, stop using this medication and get medical help right away. Precautions Before using this product, tell your doctor or pharmacist if you are allergic to naphazoline or to pheniramine; or if you have any other allergies. This product may contain inactive ingredients (such as preservatives like benzalkonium chloride), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. If you have any of the following health problems, consult your doctor or pharmacist before using this product: diabetes, glaucoma, heart problems (such as heart attack, chest pain), high blood pressure, eye infection/injury, overactive thyroid (hyperthyroidism), difficulty urinating. After you apply this drug, your vision may become temporarily blurred. Do not drive, use machinery, or do any activity that requires clear vision until you can do it safely. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: drowsiness, decreased body temperature, slow/shallow breathing. Notes Notes Avoid allergens that may be the cause of your symptoms. Some common causes of allergies include pollen, grasses, weeds, dust mites, and pet dander. Missed Dose Missed Dose If your doctor has directed you to use this medication on a regular schedule and you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Storage Storage Store at room temperature away from light. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
