

iMedix: Your Personal Health Advisor.

Amabelz

Search for medical details concerning Amabelz oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Stomach upset, nausea/vomiting, bloating, breast tenderness, headache, or weight changes may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: nausea/vomiting that doesn't stop, mental/mood changes (such as depression, memory loss), breast lumps, unusual vaginal bleeding (such as spotting, breakthrough bleeding, prolonged/recurrent bleeding), increased or new vaginal irritation/itching/odor/discharge, severe stomach/abdominal pain, yellowing eyes/skin, dark urine, swelling hands/ankles/feet, increased thirst/urination. This medication may rarely cause serious problems from blood clots (such as heart attacks, strokes, deep vein thrombosis, pulmonary embolism). Get medical help right away if you have any serious side effects, including: chest/jaw/left arm pain, unusual sweating, sudden/severe headache, weakness on one side of the body, confusion, trouble speaking, sudden vision changes (such as partial/complete blindness), pain/redness/swelling of legs, tingling/weakness/numbness in the arms/legs, trouble breathing, coughing up blood, sudden dizziness/fainting. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication contains 2 hormones: an estrogen (such as conjugated estrogen, estradiol) and a progestin (such as medroxyprogesterone, norethindrone, norgestimate). Some brands of this medication may contain progesterone instead of a progestin. Progesterone and progestin both act like the form of the natural hormone (progesterone) produced by your body. This medication is used by women to help reduce symptoms of menopause (such as hot flashes, vaginal dryness). These symptoms are caused by the body making less estrogen. The progestin or progesterone in this medication helps to reduce the risk of cancer of the uterus which can be caused by using estrogen. Women who have had their uterus removed do not need the progestin or progesterone. They should not use this combination medication. If you are using this medication to treat symptoms only in and around the vagina, products applied directly inside the vagina

should be considered before medications that are taken by mouth, absorbed through the skin, or injected. This medication may also be used by women after menopause to prevent bone loss (osteoporosis). However, there are other medications (such as raloxifene, bisphosphonates including alendronate) that are also effective in preventing bone loss and may be safer. These medications should be considered for use before estrogen/progestin treatment. How to use Amabelz Read the Patient Information Leaflet if available from your pharmacist before you start using this medication and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, usually once daily. You may take it with food or right after a meal to prevent stomach upset. The dosage is based on your medical condition and response to treatment. Take this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Follow your dosing schedule carefully. Do not increase your dose or take this medication more often or for a longer time than directed. Tell your doctor if your condition does not improve or if it worsens. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: aromatase inhibitors (such as anastrozole, exemestane, letrozole), fezolinetant, fulvestrant, ospemifene, raloxifene, tamoxifen, toremifene, tranexamic acid, certain combination products used to treat chronic hepatitis C (ombitasvir/paritaprevir/ritonavir with or without dasabuvir). This medication may interfere with certain lab tests (such as metyrapone test), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug. Precautions Before taking this medication, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: vaginal bleeding of unknown cause, certain cancers (such as breast cancer, cancer of the uterus/ovaries), blood clots, stroke, heart disease (such as heart attack), liver disease, kidney disease, family medical history (especially breast lumps, cancer, blood clots), family or personal history of a certain swelling disorder (angioedema), blood clotting disorders (such as protein C or protein S deficiency), high blood pressure, diabetes, high blood cholesterol/triglyceride levels, obesity, lupus, underactive thyroid (hypothyroidism), mineral imbalance (low or high level of calcium in the blood), a certain hormone problem (hypoparathyroidism), uterus problems (such as fibroids, endometriosis), gallbladder disease, asthma, seizures, migraine headaches, a certain blood disorder (porphyria), mental/mood disorders (such as dementia, depression). Do not smoke or use tobacco. Estrogens combined with smoking further increases your risk of stroke, blood clots, high blood pressure, and heart attack, especially in women older than 35. Tell your doctor if you just had or will be having surgery, or if you will be confined to a chair or bed for a long time (such as a long plane flight). These conditions increase your risk of getting blood clots, especially if you are using an estrogen product. You may need to stop this medication for a time or take special precautions. This medication may cause blotchy, dark areas on your face and skin (melasma). Sunlight may worsen this effect. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. If you are nearsighted or wear contact lenses, you may develop vision problems or trouble wearing your contact lenses. Contact your eye doctor if these problems occur. This medication should not be used during pregnancy. If you become pregnant or think you may be pregnant, tell your doctor right away. This medication passes into breast milk. It may reduce the quality and amount of breast milk produced. Consult your doctor before breast-feeding. Warnings Estrogens, either used alone or with a progestin, have rarely caused very serious side effects. Discuss the risks and benefits of hormone treatment with your doctor. Estrogens and progestins should not be used to prevent heart disease or dementia. In postmenopausal women, estrogen when used with a progestin can increase the risk of heart disease (such as heart attacks), stroke, serious blood clots in the lungs/legs, dementia, and cancer of the breast/ovaries. The risk for serious side effects may depend on the dose of estrogen and the length of time it is used. This medication should be used at the lowest effective dose and for the shortest amount of time. Discuss the use of this medication with your doctor regularly (for example, every 3 to 6 months) to see if you still need to use it. If you will be using this medication long-term, you should have regular complete physical exams (for example, once a year) as directed by your doctor. See also Notes section.

Side Effects

Uses

Interactions

- -
 -
 -
-

Other Details
