

iMedix: Your Personal Health Advisor.

Artesunate 110 Mg Intravenous Solution Antimalarial Drugs

Search for medical details concerning artesunate intravenous on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Unusual tiredness may occur. If this effect lasts or gets worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: easy bruising/bleeding, signs of kidney problems (such as change in amount of urine), dark urine, yellowing eyes/skin. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This medication is used to treat malaria. Malaria is an infection caused by mosquito bites received while traveling or living in regions of the world where malaria is common. Malaria parasites enter the body, and live in body tissues such as red blood cells or the liver. This medication is used to kill the malaria parasites living inside red blood cells. You will need to take a different medication (such as primaquine) to kill the malaria parasites living in the liver. Both treatments are needed for a complete cure and to avoid the return of infection (relapse). **How to use Artesunate 110 Mg Intravenous Solution Antimalarial Drugs** This medication is given by injection into a vein as directed by your doctor. The first 3 doses are usually given 12 hours apart. The following doses are usually given once daily until you are able to take an antimalarial drug by mouth. The dosage is based on your medical condition, weight, and response to treatment. Tell your doctor if your condition does not get better or if it gets worse. **Precautions** Before using artesunate, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This medication passes into breast milk in small amounts but is unlikely to harm a nursing infant. Consult your doctor before breast-feeding. **Overdose** If someone has overdosed and has serious symptoms such as passing

out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Lab and/or medical tests (such as complete blood count) should be done while you are using this medication and for 4 weeks after your last dose. Keep all medical and lab appointments. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Not applicable. This medication is given in a hospital or clinic and will not be stored at home. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details