

iMedix: Your Personal Health Advisor.

Arthricream Cream

Search for medical details concerning Arthricream topical on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Stop using this medication and tell your doctor right away if any serious side effects occur, including: blistering/peeling/redness/irritation at the application site, nausea/vomiting, ringing in the ears. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** Trolamine salicylate is used to treat minor aches and pains of the muscles/joints (such as arthritis, backache, sprains). It belongs to a class of drugs known as salicylates. **How to use Arthricream Cream** Follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. This medication is for use on the skin only. Do not get this medication in your eyes, inside your nose or mouth, or inside the vagina. If you do get the medication in those areas, flush with plenty of water. Apply the medication to the affected area no more than 3 to 4 times a day. Rub in gently and thoroughly. After applying the medication, wash your hands unless you are using this medication to treat the hands. Do not apply this medication on broken or irritated skin. Do not bandage or tightly wrap the affected area or apply heat to the area (such as using a heating pad). Doing so may increase the risk of side effects. Ask your pharmacist for more details. Tell your doctor if your condition lasts for more than 7 days, if it gets worse, or if it keeps returning. If you think you may have a serious medical problem, get medical help right away. **Precautions** Before using trolamine salicylate, tell your doctor or pharmacist if you are allergic to it; or to aspirin or other salicylates (such as salsalate); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: asthma, growths in the nose (nasal polyps). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This drug is similar to aspirin. Children and teenagers younger than 18 years should not use this drug if they have chickenpox, flu, or any undiagnosed illness or if they have

recently received a vaccine. In these cases, this drug may increase the risk of Reye's syndrome, a rare but serious illness. Tell your doctor if you are pregnant or if you plan to become pregnant. This medication may harm an unborn baby and cause problems with normal labor/delivery. It is not recommended for use in pregnancy from 20 weeks until delivery. If your doctor decides that you need to use this medication between 20 and 30 weeks of pregnancy, you should use the lowest effective dose for the shortest possible time. You should not use this medication after 30 weeks of pregnancy. Discuss the risks and benefits with your doctor. It is unknown if this product passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Keep all medical and lab appointments. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from heat. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details

