

iMedix: Your Personal Health Advisor.

Asciminib 40 Mg Tablet

Search for medical details concerning asciminib oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Muscle/bone pain, headache, tiredness, dizziness, blurred vision, nausea, vomiting, or diarrhea may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. People using this medication may have serious side effects. However, you have been prescribed this drug because your doctor has judged that the benefit to you is greater than the risk of side effects. Careful monitoring by your doctor may decrease your risk. This medication may raise your blood pressure. Check your blood pressure regularly and tell your doctor if the results are high. Your doctor may control your blood pressure with medication.

Tell your doctor right away if you have any serious side effects, including: easy bruising/bleeding, signs of anemia (such as severe tiredness, pale skin), signs of infection (such as sore throat that doesn't go away, fever, chills, cough), signs of disease of the pancreas (such as severe stomach/abdominal pain which may spread to the back, nausea/vomiting that doesn't go away). Get medical help right away if you have any very serious side effects, including: symptoms of a heart attack (such as chest/jaw/left arm pain, shortness of breath, unusual sweating), signs of a stroke (such as weakness on one side of the body, trouble speaking, sudden vision changes, confusion), signs of heart failure (such as sudden weight gain, swelling ankles/feet, trouble breathing, unusual tiredness), fast/irregular heartbeat, severe dizziness, fainting, pain/swelling/warmth in the groin/calf. Asciminib can commonly cause a rash that is usually not serious. However, you may not be able to tell it apart from a rare rash that could be a sign of a severe reaction. Get medical help right away if you develop any rash. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

How to use Asciminib 40 Mg Tablet Read the Patient Information Leaflet if available from your pharmacist before you start taking asciminib and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth without food as directed by your doctor, usually once or twice a day. Do not eat food for at least 2 hours before and at least 1 hour after taking asciminib. Swallow the tablets whole. Do not crush, chew, or break the tablets. The dosage is based on your medical condition, response to treatment, lab tests, and other medications you may be taking. Be sure to tell your doctor and pharmacist about all the products

you use (including prescription drugs, nonprescription drugs, and herbal products). Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time(s) each day. Do not increase your dose or use this drug more often or for longer than prescribed. Your condition will not improve any faster, and your risk of serious side effects will increase. Since this drug can be absorbed through the skin and lungs and may harm an unborn baby, women who are pregnant or who may become pregnant should not handle this medication or breathe the dust from the tablets. Precautions Before taking asciminib, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: disease of the pancreas, heart disease (such as heart attack, coronary artery disease, heart failure), stroke, high blood pressure. This drug may make you dizzy or cause blurred vision. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness or clear vision until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Asciminib can make you more likely to get infections or may make current infections worse. Stay away from anyone who has an infection that may easily spread (such as chickenpox, COVID-19, measles, flu). Talk to your doctor if you have been exposed to an infection or for more details. Tell your health care professional that you are using asciminib before having any immunizations/vaccinations. Avoid contact with people who have recently received live vaccines (such as flu vaccine inhaled through the nose). To lower the chance of getting cut, bruised, or injured, use caution with sharp objects like razors and nail cutters, and avoid activities such as contact sports. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Since this drug can be absorbed through the skin and lungs and may harm an unborn baby, women who are pregnant or who may become pregnant should not handle this medication or breathe the dust from the tablets. Tell your doctor if you are pregnant or plan to become pregnant. You should not become pregnant while using asciminib. Asciminib may harm an unborn baby. Your doctor should order a pregnancy test before you start this medication. Ask about reliable forms of birth control while using this medication and for 1 week after the last dose. If you become pregnant, talk to your doctor right away about the risks and benefits of this medication. It is unknown if this drug passes into breast milk. Because of the possible risk to the infant, breast-feeding is not recommended while using this drug and for 1 week after the last dose. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as blood pressure, complete blood count, lipase, amylase, heart function) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you are taking this medication once a day and miss a dose, take it as soon as you remember unless it is more than 12 hours after the time you usually take the dose. If it is more than 12 hours from the usual time of your dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. If you are taking this medication twice a day and miss a dose, take it as soon as you remember unless it is more than 6 hours after the time you usually take the dose. If it is more than 6 hours from the usual time of your dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store in the original container at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
