

# iMedix: Your Personal Health Advisor.

## Azeschew 13 Mg Iron-1 Tablet Multivitamins

---

Search for medical details concerning Azeschew oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Constipation, diarrhea, or upset stomach may occur. These effects are usually temporary and may disappear as your body adjusts to this medication. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Iron may cause your stools to turn black, an effect that is not harmful. If your doctor has prescribed this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov](http://www.fda.gov) In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is a multivitamin and iron product used to treat or prevent vitamin deficiency due to poor diet, certain illnesses, or during pregnancy. Vitamins and iron are important building blocks of the body and help keep you in good health. How to use Azeschew 13 Mg Iron-1 Tablet Multivitamins Chew this medication thoroughly and swallow, usually once daily or as directed. Follow all directions on the product package, or take as directed by your doctor. Do not take more than the recommended dosage. If you have any questions, ask your doctor or pharmacist. This medication is best taken on an empty stomach 1 hour before or 2 hours after meals. If stomach upset occurs, you may take this medication with food. Avoid taking antacids, dairy products, tea, or coffee within 2 hours before or after this medication because they will decrease its effectiveness. Take this medication regularly in order to get the most benefit from it. To help you remember, take it at the same time each day. Warnings Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children younger than 6 years. Keep this product out of reach of children. If overdose does occur, get medical help right away or call a poison control center. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug are: chloramphenicol, methyldopa. This product can decrease the absorption of other drugs such as bisphosphonates (for example, alendronate), levodopa,

penicillamine, quinolone antibiotics (for example, 500 ciprofloxacin, levofloxacin), thyroid medications (for example, levothyroxine), and tetracycline antibiotics (for example, doxycycline, minocycline). Separate your doses of these medications as far as possible from your doses of this product. Ask your doctor or pharmacist about how long you should wait between doses and for help finding a dosing schedule that will work with all your medications. If your brand of multivitamin also contains folic acid, be sure to tell your doctor or pharmacist if you take certain anti-seizure drugs (including hydantoins such as phenytoin). **Precautions** Before taking this product, tell your doctor or pharmacist if you are allergic to any of its ingredients; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before taking this medication, tell your doctor or pharmacist your medical history, especially of: iron overload disorder (such as hemochromatosis, hemosiderosis), use/abuse of alcohol, liver problems, stomach/intestinal problems (such as ulcer, colitis). If your brand of multivitamin also contains folic acid, be sure to tell your doctor or pharmacist if you have vitamin B12 deficiency (pernicious anemia) before taking it. Folic acid may affect certain lab tests for vitamin B12 deficiency without treating this anemia. Untreated vitamin B12 deficiency may result in serious nerve problems (such as peripheral neuropathy). Consult your doctor or pharmacist for details. This medicine may contain aspartame. If you have phenylketonuria (PKU) or any other condition that requires you to restrict your intake of aspartame (or phenylalanine), consult your doctor or pharmacist about using this drug safely. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant before using this medication. This medication passes into breast milk. Consult your doctor before breast-feeding.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

