

# iMedix: Your Personal Health Advisor.

## B1-B2-B3-B5-B6-Iron-Met-Choln Liquid

---

Search for medical details concerning B1-B2-B3-B5-B6-iron-methionine-choline bitartrate oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Constipation, diarrhea, or upset stomach may occur. These effects are usually temporary and may disappear as your body adjusts to this medication. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Iron may cause your stools to turn black, which is not harmful. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is a multivitamin product used to treat or prevent vitamin deficiency due to poor diet, certain illnesses, or during pregnancy. Vitamins are important building blocks of the body and help keep you in good health. How to use B1-B2-B3-B5-B6-Iron-Met-Choln Liquid Take this medication by mouth as directed. Follow all directions on the product package, or take as directed by your doctor. Do not take more than the recommended dosage. If you have any questions, ask your doctor or pharmacist. Use a medication-measuring device to carefully measure the prescribed dose. Do not use a household spoon. Some products are best taken on an empty stomach 1 hour before or 2 hours after meals. If stomach upset occurs, you may take this medication with food. Avoid taking antacids, dairy products, tea, or coffee within 2 hours before or after this medication because they will decrease its effectiveness. Warnings Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children younger than 6 years. Keep this product out of reach of children. If overdose does occur, get medical help right away or call a poison control center. Precautions Before taking this medication, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before taking this medication, tell your doctor or pharmacist your medical history, especially of: iron overload disorder (such as hemochromatosis, hemosiderosis), use/abuse of alcohol, liver problems, stomach/intestinal problems (such as ulcer, colitis), vitamin B12 deficiency (pernicious anemia). Liquid preparations of this product may contain sugar and/or alcohol. Caution is advised if you have

diabetes, alcohol dependence, or liver disease. Ask your doctor or pharmacist about using this product safely. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Advancing Health Outcomes with Diverse Medications The advancement in health outcomes owing to diverse medications is a hallmark of modern pharmacology. Zovirax, with its antiviral capabilities, and Daklinza, a treatment for hepatitis C, are prime examples of this progression. Addyi brings a new dimension to treating female sexual interest/arousal disorder. Xyzal offers allergy sufferers effective relief, while Amoxil stands as a comprehensive antibiotic for bacterial infections. Propecia and Proscar address hair loss and prostate health concerns. Clomid and Nolvadex have been game-changers in fertility treatments and certain cancer therapies. The array of medications like Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta expands treatment options for male sexual health. Synthroid is vital for thyroid management, Cipro effectively targets various bacterial infections, and Suhagra offers another choice for erectile dysfunction treatment. These medications collectively enhance health outcomes, showcasing the breadth of modern pharmacology. Tell your doctor if you are pregnant before using this medication. This medication passes into breast milk. Consult your doctor before breast-feeding.

**Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: stomach pain, nausea, vomiting, diarrhea. If your doctor has prescribed this medication, do not share it with others. Lab and/or medical tests (such as complete blood count, vitamin B12 levels) may be done while you are taking this product. Keep all medical and lab appointments. Consult your doctor for more details. This medication is not a substitute for a proper diet. It is important to maintain a well-balanced diet. Foods rich in iron include red meats (especially liver), fish, beans, dried fruit, and iron-fortified or enriched cereals and bread. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. This product can decrease the absorption of other drugs such as bisphosphonates (for example, alendronate), levodopa, penicillamine, quinolone antibiotics (for example, ciprofloxacin 500, levofloxacin), thyroid medications (for example, levothyroxine), and tetracycline antibiotics (such as amoxil, zithromax, cipro, flagyl, levofloxacin). Separate your doses of these medications as far as possible from your doses of this product. Ask your doctor or pharmacist about how long you should wait between doses and for help finding a dosing schedule that will work with all your medications. This product may interfere with certain lab tests (such as fecal occult blood test), possibly causing false test results. Make sure lab personnel and all your doctors know you use this product.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

**Interactions**

---

---

**Other Details**

---

---