

# iMedix: Your Personal Health Advisor.

## Bafiertam 95 Mg Capsule,Delayed Release Multiple Sclerosis Agents, General

---

Search for medical details concerning Bafiertam oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Flushing/warmth, redness, itching, or burning feeling of the skin may occur. Stomach/abdominal pain, diarrhea, nausea, and vomiting may also occur. Taking this medication without food may lessen stomach/abdominal side effects. These effects usually decrease or go away as your body adjusts to the medication. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication may lower your ability to fight infections. This may make you more likely to get a serious (rarely fatal) infection or make any infection you have worse. Tell your doctor right away if you have any signs of infection (such as sore throat that doesn't go away, fever, chills, cough, blisters). Tell your doctor right away if you have any serious side effects, including: symptoms of liver damage (such as nausea/vomiting that doesn't stop, loss of appetite, stomach/abdominal pain, yellowing eyes/skin, dark urine). This medication may increase your risk of getting a rare but very serious (possibly fatal) brain infection (progressive multifocal leukoencephalopathy-PML). Get medical help right away if you have any of these side effects: clumsiness, loss of coordination/balance, weakness, sudden change in your thinking (such as confusion, difficulty concentrating/memory loss), difficulty talking/walking, seizure, vision changes. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Monomethyl fumarate is used to treat multiple sclerosis (MS). It is not a cure for MS, but it may help decrease the number of relapses and prevent or delay disability. How to use Bafiertam 95 Mg Capsule,Delayed Release Multiple Sclerosis Agents, General Read the Patient Information Leaflet if available from your pharmacist before you start taking monomethyl fumarate and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, usually twice daily. To reduce your risk of side effects, your doctor may direct you to start this medication at a low dose for the first 7 days, and gradually increase your dose. Follow

your doctor's instructions carefully. Swallow the capsules whole. Do not open, crush, chew, or mix the contents with food. Doing so can release all of the drug at once, increasing the risk of side effects. If flushing occurs, ask your doctor about taking a plain (non-enteric coated) aspirin 30 minutes before taking this medication to help prevent flushing. Take this medication regularly to get the most benefit from it. To help you remember, take it at the same times each day. Tell your doctor if your condition does not get better or if it gets worse. Precautions Before taking monomethyl fumarate, tell your doctor or pharmacist if you are allergic to it; or to dimethyl fumarate; or to diroximel fumarate; or if you have any other allergies. This product may contain inactive ingredients which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: recent/past/current infections, liver disease. Monomethyl fumarate can make you more likely to get infections or may make current infections worse. Stay away from anyone who has an infection that may easily spread (such as chickenpox, COVID-19, measles, flu). Talk to your doctor if you have been exposed to an infection or for more details. Tell your health care professional that you are using monomethyl fumarate before having any immunizations/vaccinations. Avoid contact with people who have recently received live vaccines (such as flu vaccine inhaled through the nose). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug are: other drugs that weaken the immune system/increase the risk of infection (such as natalizumab, rituximab). Monomethyl fumarate is very similar to dimethyl fumarate and diroximel fumarate. Do not use medications containing dimethyl fumarate or diroximel fumarate while using monomethyl fumarate. Does Bafiertam 95 Mg Capsule,Delayed Release Multiple Sclerosis Agents, General interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as complete blood counts, liver function) should be done before you start taking this medication and while you are taking it. Keep all medical and lab appointments. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store the unopened bottle in the refrigerator away from light and moisture. After opening, keep this medication in the original container and store at room temperature. Do not store in the bathroom. Keep all medications away from children and pets. After the bottle is opened, throw away any unused medication after 3 months. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

## Side Effects

---

## Uses

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---