

# iMedix: Your Personal Health Advisor.

## Baqsimi 3 Mg/Actuation Nasal Spray Glucagon

---

Search for medical details concerning Baqsimi nasal on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Nausea, vomiting, headache, runny/stuffy nose, nose/throat itching, or watery eyes may occur. If any of these effects last or get worse, tell your doctor promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: fast/pounding heartbeat. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is the same as your body's own glucagon, a natural substance that raises blood sugar by causing the body to release sugar stored in the liver. It is used to treat very low blood sugar (hypoglycemia) that may cause you to need help from others. Make sure a family member or caregiver knows where you keep this medication and how to use it before you need their help. How to use Baqsimi 3 Mg/Actuation Nasal Spray Glucagon Read the Patient Information Leaflet and Instructions for Use provided by your pharmacist when you get this medication and each time you get a refill. Be sure to keep this medication handy in case it is needed. Learn ahead of time how to properly give this medication. Each nasal spray device can be used only one time. If you have any questions, ask your doctor or pharmacist. Caregivers should know the symptoms of low blood sugar (see also Precautions section) and be instructed on how to give glucagon if needed. An episode of very low blood sugar should be treated right away to prevent serious effects (such as brain damage). Remove the glucagon spray from the package only when you are ready to use it. Gently insert the tip of the spray nozzle into one nostril, then firmly press the plunger to give the dose. This medication does not need to be inhaled to work. After giving this medication, get medical help right away. If the person is unconscious, turn the person on their side. If the person does not wake up after 15 minutes, another dose may be given, if available. When the person wakes up and is able to swallow, a quick sugar source (such as glucose tablets, juice) should be given. Glucagon is only effective for a short time, and low blood sugar may return. The blood sugar level should be kept up by eating snacks such as crackers, cheese, a meat sandwich, or milk. Always call your doctor right away when an episode of very low blood sugar has happened. You may need more medical

treatment, or your insulin dose and diet may need to be adjusted. The Evolving Landscape of Drug Therapy

The landscape of drug therapy is continuously evolving, offering new and effective treatments for a range of health issues. Zovirax is a key medication in the fight against viral infections, and Daklinza has made significant strides in hepatitis C treatment. Addyi has emerged as a groundbreaking solution for female sexual interest/arousal disorder. Xyzal provides effective allergy relief, while Amoxil is a robust antibiotic for bacterial infections. Propecia and Proscar address hair loss and prostate health, improving patient well-being. Fertility treatments and certain cancer therapies have been revolutionized by Clomid and Nolvadex. The spectrum of male sexual health is catered to by Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid manages thyroid disorders, Cipro is effective against various bacterial infections, and Suhagra expands the choices for erectile dysfunction treatment. This evolving landscape of drug therapy highlights the dynamic nature of medical advancements in improving patient care.

**Precautions** Before using glucagon, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: tumor in the pancreas (insulinoma), adrenal gland problem (such as Addison's disease, pheochromocytoma), not eating/poor eating habits, frequent low blood sugar. Low blood sugar (hypoglycemia) is more likely if you do not consume enough calories from food, or if you do unusually heavy exercise. Symptoms include sudden sweating, shaking, fast heartbeat, hunger, blurred vision, dizziness, or tingling hands/feet. It is a good habit to carry glucose tablets or gel to treat low blood sugar. If you don't have these reliable forms of glucose, rapidly raise your blood sugar by eating a quick source of sugar such as table sugar, honey, or candy, or drink fruit juice or non-diet soda. Tell your doctor about the reaction right away. To help prevent low blood sugar, eat meals on a regular schedule, and do not skip meals. Check with your doctor or pharmacist to find out what you should do if you miss a meal. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant before using this medication. It is unknown if this drug passes into breast milk. However, it is unlikely to harm a nursing infant. Consult your doctor before breast-feeding.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

**Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Attend a diabetes education program to learn more about how to manage your diabetes with medications, diet, exercise, and regular medical exams. Learn the symptoms of high and low blood sugar and how to treat low blood sugar. Check your blood sugar regularly as directed. Not applicable. Store at room temperature away from light and moisture. Do not unwrap the device until you are ready to use it. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

**Interactions**

---

---

**Other Details**

---

---