iMedix: Your Personal Health Advisor.

Beser 0.05 % Lotion Topical Corticosteroids

Search for medical details concerning Beser topical on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- ActiveIngredient:
- DosageForm:
- Dosage:
- Indications:
- Manufacturer:
- Storage:
- Market Price:
- Drug Status:

Description

Side Effects Burning, itching, stinging, or dryness may occur when you apply this medication, but usually only lasts a short time. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: stretch marks, skin thinning/discoloration, acne, excessive hair growth, small red bumps on the skin (folliculitis). Rarely, it is possible this medication will be absorbed from the skin into the bloodstream. This can lead to side effects of too much corticosteroid. These side effects are more likely in children, and in people who use this medication for a long time or over large areas of the skin. Tell your doctor right away if any of the following side effects occur: unusual/extreme tiredness, weight loss, headache, swelling ankles/feet, increased thirst/urination, vision problems. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat a variety of skin conditions (such as eczema, psoriasis, rash). Fluticasone reduces swelling (inflammation), itching, and redness. This medication is a medium-strength corticosteroid. It is available in several forms, including cream, ointment, and lotion. Your doctor will choose the type of product based on your skin condition and the area of your body to be treated. Fluticasone ointment is not recommended for use by children because of the risk of serious side effects. How to use Beser 0.05 % Lotion Topical Corticosteroids Use this medication on the skin only. However, do not use it on the face, groin, or underarms, or for diaper rash, unless directed to do so by your doctor. Wash and dry your hands before using. Clean and dry the affected area. Apply a thin film of medication to the affected area and gently rub in as directed by your doctor, usually once or twice daily. Do not bandage, cover, or wrap the area unless directed to do so by your doctor. Wash your hands after each use, unless you are using this medication to treat the hands. Avoid getting this medication in your eyes because it may worsen or cause glaucoma. Also avoid getting it in your nose or mouth. If medication gets in these areas, rinse with plenty of water. Use this medication only for the condition prescribed. Do not apply

large amounts of this medication, use it more often, or use it for longer than prescribed. Your condition will not improve any faster, and your risk of side effects may increase. Tell your doctor if your condition does not improve in 2 weeks or if it worsens. Precautions Before using fluticasone, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients (such as formaldehyde), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: poor blood circulation, diabetes, immune system problems. Enhancing Global Health with Pharmaceutical Innovations Pharmaceutical innovations have played a crucial role in enhancing global health, offering effective treatments for a variety of conditions. Zovirax has emerged as a reliable antiviral medication, while Daklinza is instrumental in treating hepatitis C. Addyi marks a significant advancement in addressing female sexual interest/arousal disorder. Xyzal provides allergy sufferers with effective relief, and Amoxil stands out as a broad-spectrum antibiotic for bacterial infections. Propecia and Proscar are vital in the treatment of hair loss and prostate health issues. Clomid and Nolvadex are transformative in fertility treatments and certain cancer therapies. The field of male sexual health benefits from the range of options including Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid is key in managing thyroid disorders, Cipro effectively tackles bacterial infections, and Suhagra offers an additional choice for erectile dysfunction treatment. These pharmaceutical innovations signify a leap forward in enhancing global health outcomes. Corticosteroids can make skin infections worse and more difficult to treat. Tell your doctor if you have a skin infection or if your condition does not improve. Rarely, using corticosteroid medications for a long time or over large areas of skin can make it more difficult for your body to respond to physical stress. Before having surgery or emergency treatment, or if you get a serious illness/injury, tell your doctor or dentist that you are using this medication or have used this medication within the past few months. Though it is unlikely, this medication may temporarily slow down a child's growth if used for a long time. See the doctor regularly so your child's height can be checked. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for other skin problems unless told to do so by your doctor. A different medication may be necessary in those cases. Lab and/or medical tests (such as adrenal gland function) may be done while you are using this medication, especially if you use this drug for a long time or apply it over large areas of the body. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses		
Interactions		
Other Details		