

iMedix: Your Personal Health Advisor.

Bicalutamide

Search for medical details concerning bicalutamide oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Hot flashes, breast swelling/tenderness/pain, nausea, vomiting, constipation, diarrhea, stomach upset, weight changes, headache, trouble sleeping, drowsiness, or dizziness may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: increased thirst/urination, decreased sexual ability/desire, mental/mood changes (such as anxiety, depression), swelling hands/ankles/feet, shortness of breath, increased coughing, numb/tingling skin, bone/joint pain, unusual tiredness, muscle weakness, painful/bloody urination, signs of infection (such as sore throat that doesn't go away, fever, chills). Get medical help right away if you have any very serious side effects, including: chest/jaw/left arm pain. This medication may rarely cause serious (possibly fatal) liver disease. Get medical help right away if any of these side effects occur: nausea/vomiting that doesn't stop, loss of appetite, yellowing eyes/skin, stomach/abdominal pain, dark urine. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Bicalutamide is used to treat prostate cancer. This medication works by blocking the action of hormones in the prostate, slowing the growth of cancer cells. This medication should not be used in women and children. How to use Bicalutamide Read the Patient Information Leaflet if available from your pharmacist before you start taking bicalutamide and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, usually once a day. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Do not increase your dose or use this drug more often or for longer than prescribed. Your condition will not improve any faster, and your risk of serious side effects will increase. Since this drug can be absorbed through the skin and lungs and may harm an unborn baby, women who are pregnant or who may become pregnant should not handle this medication or breathe the dust from the tablets. Precautions Before taking bicalutamide, tell your doctor or pharmacist if you are allergic to it; or if you have

any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, diabetes, heart disease. This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). This medication may make you more sensitive to the sun. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned or have skin blisters/redness. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication should not be used in women, especially during pregnancy or breast-feeding. It may harm an unborn or breast-feeding baby. Consult your doctor for more details. Male patients and their female partners should use reliable forms of birth control during and for 130 days after treatment with bicalutamide since this medication may damage sperm. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug are: certain "blood thinners" (such as warfarin). Does Bicalutamide interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as PSA levels, liver function, blood sugar, complete blood count) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details
