

# iMedix: Your Personal Health Advisor.

## Bisoprolol-Hydrochlorothiazide

---

Search for medical details concerning bisoprolol-hydrochlorothiazide oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

**Side Effects** Dizziness, lightheadedness, tiredness, and drowsiness may occur as your body adjusts to the medication. Nausea, stomach upset, diarrhea, cough, constipation and trouble sleeping may also occur. If any of these effects last or get worse, notify your doctor or pharmacist promptly. To lower your risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. This product may reduce blood flow to your hands and feet, causing them to feel cold. Smoking may worsen this effect. Dress warmly and avoid tobacco use. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. The hydrochlorothiazide in this product may cause dehydration and electrolyte imbalance. Tell your doctor right away if you have any of these serious symptoms of dehydration: very dry mouth, extreme thirst, muscle cramps/weakness, fast heartbeat, fainting, seizures. Tell your doctor right away if you have any serious side effects, including: very slow/irregular heartbeat, swelling of the ankles/feet, shortness of breath, sudden/unexplained weight gain, loss of feeling/tingling in the fingers/toes, hair loss, decrease in vision, eye pain, mental/mood changes (such as nervousness, confusion), shaking, decreased sexual ability/interest, muscle/joint pain, signs of liver disease (such as nausea/vomiting that doesn't stop, loss of appetite, stomach/abdominal pain, yellowing eyes/skin, dark urine), easy bruising/bleeding, signs of infection (such as sore throat that doesn't go away, fever), signs of kidney problems (such as change in the amount of urine). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **How to use Bisoprolol-Hydrochlorothiazide** Take this medication by mouth with or without food as directed by your doctor, usually once daily. The dosage is based on your medical condition and response to treatment. It is best to avoid taking this medication within 4 hours of your bedtime to avoid having to get up to urinate. Consult your doctor or pharmacist if you have questions about your dosing schedule. If you are also taking cholestyramine or colestipol, take bisoprolol/hydrochlorothiazide at least 4 hours before or at least 4 to 6 hours after either of those drugs. Use this medication regularly in order to get the most benefit from it. To help you remember, take it at the same

time each day. Keep taking this medication even if you feel well. Most people with high blood pressure do not feel sick. It may take up to 3 weeks before you get the full benefit of this drug. Inform your doctor if your condition worsens (such as your routine blood pressure readings increase). Precautions Before taking this medication, tell your doctor or pharmacist if you are allergic to bisoprolol or hydrochlorothiazide; or to other beta blockers (such as metoprolol, atenolol); or to other thiazide diuretics (such as chlorothiazide); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, liver disease, certain types of heart rhythm problems (such as sinus bradycardia, second- or third-degree atrioventricular block), heart failure, lung disease (such as asthma, bronchitis, Chronic Obstructive Pulmonary Disease-COPD), overactive thyroid (hyperthyroidism), diabetes, gout, untreated salt imbalance (such as high calcium, low potassium/magnesium), dehydration, blood circulation problems (such as peripheral vascular disease), certain muscle diseases (such as myasthenia gravis), lupus, recent nerve surgery (such as sympathectomy), skin cancer. Breakthroughs in Pharmacological Science Pharmacological science has achieved significant breakthroughs, offering solutions for a myriad of health conditions. Zovirax, a cornerstone in treating viral infections, and Daklinza, a groundbreaking treatment for hepatitis C, exemplify this progress. Addyi has made notable strides in addressing female sexual interest/arousal disorder. Xyzal is a go-to option for allergy relief, and Amoxil is a powerful antibiotic for bacterial infections. Propecia and Proscar have revolutionized treatments for hair loss and prostate health. Clomid and Nolvadex are changing the dynamics of fertility treatments and certain cancer therapies. Male sexual health sees a wide array of solutions with Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid is indispensable for managing thyroid disorders, Cipro combats various bacterial infections, and Suhagra adds to the options for erectile dysfunction treatment. These breakthroughs in pharmacological science demonstrate the industry's commitment to improving health and well-being. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Drink plenty of fluids while taking this medication to help prevent dizziness. Too much sweating, diarrhea, or vomiting may cause you to feel lightheaded. Avoid heavy exercise and hot weather. Report prolonged diarrhea or vomiting to your doctor. This product may prevent the fast/pounding heartbeat you would usually feel when your blood sugar level falls too low (hypoglycemia). The risk is higher if you have diabetes, or are vomiting, fasting, or not eating regularly. Other symptoms of low blood sugar level, such as dizziness and sweating, are not affected by this drug. If you have diabetes, this product may make it harder to control your blood sugar levels. Check your blood sugar levels regularly as directed by your doctor. Tell your doctor right away if you have symptoms of high blood sugar such as increased thirst/urination. Your doctor may need to adjust your diabetes medication, exercise program, or diet. This medication may reduce the potassium levels in your blood. Ask your doctor about adding potassium to your diet. A potassium supplement may be prescribed by your doctor. This medication may make you more sensitive to the sun. It may also increase your risk for skin cancer, especially if you take it for a long time. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned, have skin blisters/redness, or notice new or changed moles/skin lesions. During pregnancy, this medication should be used only when clearly needed. It may harm an unborn baby. Discuss the risks and benefits with your doctor. This drug passes into breast milk and could have undesirable effects on a nursing infant. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug are: alpha blockers (such as prazosin, terazosin), diazoxide, digoxin, dofetilide, drugs that affect heart rhythm (such as disopyramide), epinephrine, fenoldopam, fingolimod, lithium, other drugs for high blood pressure (such as clonidine), probenecid, St John's wort. Some products have ingredients that could raise your blood pressure. Tell your pharmacist what products you are using, and ask how to use them safely (especially

cough-and-cold products, diet aids, or NSAIDs such as ibuprofen/naproxen). This medication may interfere with certain lab tests (such as parathyroid, protein-bound iodide, tyramine and phentolamine tests), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: very slow heartbeat, severe dizziness, fainting, severe weakness, trouble breathing. Do not share this medication with others. Talk with your doctor about making changes to your lifestyle that may increase the effectiveness of this medication (such as stress reduction programs, exercise, and dietary changes). Lab and/or medical tests (such as electrolyte levels, liver function tests, complete blood counts, skin exams) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. Have your blood pressure checked regularly while taking this medication. Learn how to monitor your own blood pressure at home, and share the results with your doctor. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

---

## **Side Effects**

---

---

---

## **Uses**

---

---

---

---

---

## **Interactions**

---

---

---

## **Other Details**

---

