

iMedix: Your Personal Health Advisor.

CALMOL 4 Suppository, Rectal

Search for medical details concerning Calmol-4 rectal on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects This medication usually does not have side effects when used as directed. If you have any unusual effects, contact your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

Uses This medication is used to temporarily relieve burning, pain, and itching caused by hemorrhoids. This product contains ingredients such as cocoa butter, starch, or zinc oxide that form a protective barrier to prevent too much irritating contact with stool. This barrier helps protect inflamed, irritated skin and helps make bowel movements less painful. Do not use this product in children 12 years or younger without talking with the doctor.

How to use CALMOL 4 Suppository, Rectal Follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. Use this product in the rectum, usually up to 6 times a day, or after each bowel movement, or as directed by your doctor. The dosage is based on your medical condition and response to treatment. Do not use more often or for a longer time than directed. Before use, clean the area with mild soap and water, rinse well, and pat dry. If the suppository is too soft to insert, place it in cold water or refrigerate it for 30 minutes before removing the wrapper. To use suppositories, unwrap the suppository. Lie on your left side with the right knee bent. Carefully push the suppository (pointed side first) just inside the anus with your finger. Do not push more than an inch or two inside the anus. Do not use this medication if hemorrhoids are bleeding or the rectal area is raw and irritated. Tell your doctor if your condition does not improve after 7 days or if bleeding/worsening pain occurs. If you think you may have a serious medical problem, get medical help right away.

Precautions Before using this product, tell your doctor or pharmacist if you are allergic to any of its ingredients, or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history. Before having surgery, tell your doctor or dentist about all the products you

use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. However, it is unlikely to pass into breast milk or to harm a nursing infant. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Does CALMOL 4 Suppository, Rectal interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. To maintain normal bowel function and reduce the risk of hemorrhoids, eat a proper diet high in fiber, drink 6 to 8 glasses of water each day, and exercise regularly. If you are using this product on a regular schedule and miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Different brands of this medication may have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details

