

# iMedix: Your Personal Health Advisor.

## CARISOPRODOL COMPOUND

---

Search for medical details concerning carisoprodol-aspirin oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Drowsiness, dizziness, headache, stomach upset, nausea/vomiting, or heartburn may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: nausea/vomiting that doesn't stop, easy bruising/bleeding, dark urine, yellowing eyes/skin, hearing problems (such as ringing in the ears, hearing loss), signs of kidney problems (such as change in the amount of urine), fast/pounding heartbeat, unusual tiredness, confusion. This drug may rarely cause serious (rarely fatal) bleeding from the stomach or intestines. If you notice any of the following unlikely but serious side effects, stop taking this medication and consult your doctor or pharmacist right away: stomach/abdominal pain that doesn't go away, black stool, vomit that looks like coffee grounds. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This product is used short-term to treat muscle pain and discomfort. It is usually used along with rest, physical therapy, and other treatments. Carisoprodol helps to relax the muscles. Aspirin helps to decrease pain and swelling. How to use CARISOPRODOL COMPOUND Take this medication by mouth with or without food as directed by your doctor. Do not increase your dose or use this drug more often or for longer than prescribed. Your condition will not improve any faster, and your risk of side effects will increase. Take this medication with a full glass of water (8 ounces/240 milliliters) unless your doctor directs you otherwise. Do not lie down for at least 10 minutes after taking this medication. If you have stomach upset while taking this medication, take it with food or milk. The dosage is based on your medical condition and response to treatment. This medication should only be used short-term (for 3 weeks or less) unless directed by your doctor. Use the smallest effective dose. Pain medications work best if they are used as the first signs of pain occur. If you wait until the symptoms have worsened, the medication may not work as well. If you suddenly stop using this medication, you may have withdrawal symptoms (such as stomach cramps, trouble sleeping, headache,

nausea). To help prevent withdrawal, your doctor may lower your dose slowly. Withdrawal is more likely if you have used this medication for a long time or in high doses. Tell your doctor or pharmacist right away if you have withdrawal. Though it helps many people, this medication may sometimes cause addiction. This risk may be higher if you have a substance use disorder (such as overuse of or addiction to drugs/alcohol). Take this medication exactly as prescribed to lower the risk of addiction. Ask your doctor or pharmacist for more details. Tell your doctor if your condition lasts after 2 to 3 weeks or if it gets worse.

**Precautions** Before taking this medication, tell your doctor or pharmacist if you are allergic to carisoprodol or aspirin; or to meprobamate, tybamate, or mebutamate; or to nonsteroidal anti-inflammatory drugs-NSAIDs (such as ibuprofen); or to other salicylates (such as choline salicylate); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: a certain blood disorder (acute intermittent porphyria), asthma (including a history of worsening breathing with runny/stuffy nose after taking aspirin or other NSAIDs), growths in the nose (nasal polyps), stomach/intestine/esophagus problems (such as bleeding, heartburn, ulcers), bleeding/clotting disorders (such as hemophilia, von Willebrand disease, thrombocytopenia), personal or family history of a substance use disorder (such as overuse of or addiction to drugs/alcohol), kidney disease, liver disease, seizure, certain enzyme deficiencies (such as G6PD or pyruvate kinase deficiency), gout. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). This medicine may cause stomach bleeding. Daily use of alcohol and tobacco, especially when combined with this medicine, may increase your risk for stomach bleeding. Avoid alcohol and stop smoking. Consult your doctor or pharmacist for more information. This medication contains aspirin. Children and teenagers less than 18 years old should not take aspirin if they have chickenpox, flu, or any undiagnosed illness or if they have recently received a vaccine. In these cases, taking aspirin increases the risk of Reye's syndrome, a rare but serious illness. Older adults may be more sensitive to the side effects of this drug, especially drowsiness, confusion, stomach/intestinal bleeding, and ulcers. Drowsiness and confusion can increase the risk of falling. Before using this medication, women of childbearing age should talk with their doctor(s) about the benefits and risks. Tell your doctor if you are pregnant or if you plan to become pregnant. This medication may harm an unborn baby and cause problems with normal labor/delivery. It is not recommended for use in pregnancy from 20 weeks until delivery. If your doctor decides that you need to use this medication between 20 and 30 weeks of pregnancy, you should use the lowest effective dose for the shortest possible time. You should not use this medication after 30 weeks of pregnancy. This medication passes into breast milk and may have undesirable effects on a nursing infant. Breast-feeding while using this drug is not recommended. Consult your doctor before breast-feeding.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: corticosteroids (such as prednisone), mifepristone. This medication may increase the risk of bleeding when taken with other drugs that also may cause bleeding. Examples include anti-platelet drugs such as clopidogrel/ticagrelor, "blood thinners" such as dabigatran/enoxaparin/warfarin, among others. Check all prescription and nonprescription medicine labels carefully since many medications contain pain relievers/fever reducers (aspirin, NSAIDs such as ibuprofen or naproxen). These drugs if taken together with this medication may increase your risk of side effects. However, if your doctor has directed you to take low-dose aspirin to prevent heart attack or stroke (usually 81-162 milligrams a day), you should continue taking the aspirin unless your doctor instructs you otherwise. Ask your doctor or pharmacist for more details. Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), other muscle relaxants (such as cyclobenzaprine, methocarbamol), or antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause

drowsiness. Ask your pharmacist about using those products safely. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: ringing in ears, severe drowsiness/dizziness, seizures, slow/shallow breathing, mental/mood changes (such as confusion, hallucinations), inability to move your legs/arms, shaky/unsteady movement, vision changes (such as blurred vision). Do not share this medication with others. Sharing it is against the law. This medication has been prescribed for your current condition only. Do not use it later for another condition unless your doctor directs you to do so. A different medication may be necessary in that case. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## Side Effects

---

## Uses

---

---

---

---

---

---

## Interactions

---

---

## Other Details

---

---