iMedix: Your Personal Health Advisor.

Cefotetan 1 Gram Solution For Injection

Search for medical details concerning cefotetan injection on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- ActiveIngredient:
- DosageForm:
- Dosage:
- Indications:
- Manufacturer:
- Storage:
- Market Price:
- Drug Status:

Description

Side Effects Swelling, redness, pain, or soreness at the injection site may occur. This medication may also rarely cause loss of appetite, nausea, vomiting, diarrhea, or headache. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: dark urine, easy bruising/bleeding, fast/pounding/irregular heartbeat, seizures, unusual weakness, yellowing eyes/skin, mental/mood changes (such as confusion). This medication may rarely cause a severe intestinal condition due to a bacteria called C. difficile. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. Use of this medication for prolonged or repeated periods may result in oral thrush or a new vaginal yeast infection. Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge, or other new symptoms. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Cefotetan is an antibiotic used to treat a wide variety of bacterial infections. It may also be used before and during certain surgeries to help prevent infection. This medication is known as a cephalosporin antibiotic. It works by stopping the growth of bacteria. This antibiotic treats only bacterial infections. It will not work for virus infections (such as common cold, flu). Using any antibiotic when it is not needed can cause it to not work for future infections. How to use Cefotetan 1 Gram Solution For Injection Read and learn all preparation and usage instructions supplied by the manufacturer. Follow all instructions for proper mixing with the correct IV fluids. Consult your pharmacist for details. Before using, check the product visually for particles or discoloration. If either is present, do not use the liquid. If you are using the frozen pre-mixed solution, thaw the container at room temperature or in the refrigerator. If the bag is thawed in the refrigerator, let it sit at room temperature at least 1 hour before using.

Do not thaw by putting in a water bath or microwaving. After thawing, shake well and squeeze the container to check for leaks. Discard solution if the container leaks. Do not re-freeze the solution after thawing. This medication is given by injection into a muscle or vein as directed by your doctor. For the best effect, use this antibiotic at evenly spaced times. To help you remember, use this medication at the same time(s) every day. The dosage is based on your medical condition and response to treatment. Continue to use this medication until the full prescribed treatment period is finished, even if symptoms disappear after a few days. Stopping the medication too early may result in a return of the infection. Learn how to store and discard needles and medical supplies safely. Consult your pharmacist for details. Precautions Before using cefotetan, tell your doctor or pharmacist if you are allergic to it; or to other cephalosporins; or to penicillins; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, liver disease, stomach/intestinal diseases (such as colitis), use/abuse of alcohol. Avoid alcoholic beverages while taking this medication and for at least 3 days after finishing this medicine. Drinking alcohol may result in severe stomach upset/cramps, nausea, vomiting, headache, and flushing. Cefotetan may cause live bacterial vaccines (such as typhoid vaccine) to not work well. Tell your health care professional that you are using cefotetan before having any immunizations/vaccinations. Tell your doctor if you are pregnant before using this medication. This medication passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with these drugs are: aminoglycoside antibiotics (such as amoxil, zithromax, cipro, flagyl, levofloxacin), "blood thinners" (such as warfarin), chloramphenicol. This medication may interfere with certain lab tests (such as as urine glucose tests, serum or urine creatinine tests), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug. Does Cefotetan 1 Gram Solution For Injection interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: trouble breathing, seizures. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to. Lab and/or medical tests (such as complete blood count, kidney function) may be done while you are using this product. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Before mixing, store at room temperature away from light and moisture. Do not refrigerate or freeze the dry powder. Different brands/packaging of this medication may have different storage requirements. Read the package labeling or ask your pharmacist for the storage requirements before and after mixing for the product you are using. Do not store in the bathroom. Store the pre-mixed solution containers in the freezer at or below -4 degrees F (-20 degrees C). (For thawing instructions, see How to Use section.) If stored at room temperature, use/discard the thawed solution within 24 hours. Consult the package instructions or your pharmacist for other storage details. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

| Uses | | |
|---------------|--|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| Interactions | | |
| | | |
| Other Details | | |
| | | |