iMedix: Your Personal Health Advisor.

Cefoxitin Vial

Search for medical details concerning cefoxitin intravenous on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- ActiveIngredient:
- DosageForm:
- Dosage:
- Indications:
- Manufacturer:
- Storage:
- Market Price:
- Drug Status:

Description

Side Effects Swelling, redness, pain, or soreness at the injection site may occur. This medication may also rarely cause loss of appetite, nausea, vomiting, diarrhea, or headache. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: dark urine, easy bruising/bleeding, fast/pounding/irregular heartbeat, seizures, unusual weakness, yellowing eyes/skin, mental/mood changes (such as confusion). This medication may rarely cause a severe intestinal condition due to a bacteria called C. difficile. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. Use of this medication for prolonged or repeated periods may result in oral thrush or a new vaginal yeast infection. Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge, or other new symptoms. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Cefoxitin is an antibiotic used to treat a wide variety of bacterial infections. It may also be used before and during certain surgeries to help prevent infection. This medication is known as a cephalosporin antibiotic. It works by stopping the growth of bacteria. How to use Cefoxitin Vial This medication is given by injection into a muscle or vein as directed by your doctor. The dosage is based on your medical condition and response to treatment. If you are giving this medication to yourself at home, learn all preparation and usage instructions from your health care professional. Before using, check this product visually for particles or discoloration. If either is present, do not use the liquid. Learn how to store and discard medical supplies safely. For the best effect, use this antibiotic at evenly spaced times. To help you remember, use this medication at the same time(s) every day. Continue to use this medication until the full prescribed treatment period is finished, even if symptoms disappear after a few days. Stopping the medication too early may result in a return of the infection. Tell your doctor if your condition lasts or gets worse. Precautions Before using cefoxitin, tell your doctor or pharmacist if you are allergic to it; or to other cephalosporins; or to penicillins; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, liver disease, stomach/intestinal diseases (such as colitis). Cefoxitin may cause live bacterial vaccines (such as typhoid vaccine) to not work well. Tell your health care professional that you are using cefoxitin before having any immunizations/vaccinations. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant before using this medication. This medication passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. This medication may interfere with certain lab tests (such as Coombs' test, urine glucose tests, serum or urine creatinine tests), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug. Does Cefoxitin Vial interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: trouble breathing, seizures. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to. Lab and/or medical tests (such as complete blood count, kidney function) may be done while you are using this product. Keep all medical and lab appointments. Consult your doctor for more details. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Do not double the dose to catch up. Consult the product instructions and your pharmacist for storage details. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects			
Uses			

Interactions			
Other Details			