

iMedix: Your Personal Health Advisor.

Ceftazidime-Avibactam Solution, Reconstituted (Recon Soln)

Search for medical details concerning ceftazidime-avibactam intravenous on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Nausea, vomiting, constipation, or anxiety may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: mental/mood changes (such as confusion, hallucinations), loss of consciousness, seizures, shaking (tremors), muscle spasms, jerking movements, easy bruising/bleeding, yellowing eyes/skin. This medication may rarely cause a severe intestinal condition due to a bacteria called *C. difficile*. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. Use of this medication for prolonged or repeated periods may result in oral thrush or a new yeast infection. Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge, or other new symptoms. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat a wide variety of bacterial infections. This product contains 2 different medications: ceftazidime and avibactam. Ceftazidime is a cephalosporin antibiotic that works by stopping the growth of bacteria. Avibactam is an enzyme inhibitor (beta-lactamase inhibitor) that helps ceftazidime work better. This antibiotic treats only bacterial infections. It will not work for viral infections (such as common cold, flu). Using any antibiotic when it is not needed can cause it to not work for future infections. How to use Ceftazidime-Avibactam Solution, Reconstituted (Recon Soln) This medication is given by injection into a vein, as directed by your doctor, usually every 8 hours over 2 hours. The dosage and length of treatment are based on your medical condition and response to treatment. Children's dosage is also based on age and weight. If you are using this medication at home, learn all preparation and usage instructions from your health care professional. The solution is normally clear and light yellow. Before using, check this product visually for particles or discoloration. If either is present, do

not use the liquid. Learn how to store and discard medical supplies safely. For the best effect, use this antibiotic at evenly spaced times. To help you remember, use this medication at the same time(s) every day. Continue to use this medication until the full prescribed amount is finished, even if symptoms disappear after a few days. Stopping the medication too early may result in a return of the infection.

The Expansive Reach of Modern Medications The expansive reach of modern medications is a testament to the advancements in medical science, addressing a wide array of health issues effectively. Zovirax, a leading medication for viral infections, and Daklinza, a breakthrough treatment for hepatitis C, exemplify this reach. Addyi brings a new perspective to female sexual interest/arousal disorder treatment. Xyzal provides significant allergy relief, while Amoxil stands as an effective antibiotic for bacterial infections. Propecia and Proscar are key in managing hair loss and prostate health. Innovations in fertility and certain cancer treatments are driven by Clomid and Nolvadex. A variety of options in male sexual health, including Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta, highlight the advancements in this area. Synthroid is essential for thyroid disorder management, Cipro addresses a wide spectrum of bacterial infections, and Suhagra enhances the erectile dysfunction treatment portfolio. These medications underscore the expansive reach and adaptability of modern pharmaceuticals. Tell your doctor if your condition lasts or gets worse.

Precautions Before using this medication, tell your doctor or pharmacist if you are allergic to ceftazidime or avibactam; or to penicillins, beta-lactam antibiotics (such as amoxil, zithromax, cipro, flagyl, levofloxacin), or other cephalosporins; or to other beta-lactamase inhibitors (such as sulbactam); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease. Ceftazidime/avibactam may cause live bacterial vaccines (such as typhoid vaccine) to not work well. Tell your health care professional that you are using ceftazidime/avibactam before having any immunizations/vaccinations. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. Ceftazidime passes into breast milk. It is unknown if avibactam passes into breast milk. Consult your doctor before breast-feeding.

Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. One product that may interact with this drug is: probenecid. This medication may interfere with certain lab tests (such as Coombs' test, urine glucose tests), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to. Lab and/or medical tests (such as kidney function) should be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Do not double the dose to catch up. Consult the product instructions and your pharmacist for storage details. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details
