

iMedix: Your Personal Health Advisor.

Centrum Silver Women

Search for medical details concerning Centrum Silver Women oral on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Constipation, diarrhea, or upset stomach may occur. These effects are usually temporary and may disappear as your body adjusts to this medication. If any of these effects last or get worse, contact your doctor or pharmacist promptly. Iron may cause your stools to turn black, an effect that is not harmful. If your doctor has prescribed this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is a multivitamin and iron product used to treat or prevent vitamin deficiency due to poor diet, certain illnesses, or during pregnancy. Vitamins and iron are important building blocks of the body and help keep you in good health. How to use Centrum Silver Women Take this medication by mouth, usually once daily or as directed. Follow all directions on the product package, or take as directed by your doctor. Do not take more than the recommended dosage. If you have any questions, ask your doctor or pharmacist. This medication is best taken on an empty stomach 1 hour before or 2 hours after meals. Take with a full glass of water (8 ounces or 240 milliliters) unless your doctor directs you otherwise. If stomach upset occurs, you may take this medication with food. Avoid taking antacids, dairy products, tea, or coffee within 2 hours before or after this medication because they may decrease its effectiveness. Do not lie down for at least 10 minutes after taking the tablets or capsules. Consult your doctor or pharmacist for details for your particular brand. If you are taking the delayed-release form or extended-release capsules, swallow them whole. Do not crush or chew delayed-release or extended-release products. Doing so can release all of the drug at once, increasing the risk of side effects and decreasing absorption. Also, do not split extended-release tablets unless they have a score line and your doctor or pharmacist tells you to do so. Swallow the whole or split tablet without crushing or chewing. Take this medication regularly in order to get the most benefit from it. To help you remember, take it at the same time each day. Warnings Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children younger than 6 years. Keep this product out of reach of children. If overdose does occur, get medical

help right away or call a poison control center. Precautions Before taking this product, tell your doctor or pharmacist if you are allergic to any of its ingredients; or if you have any other allergies. This product may contain inactive ingredients (such as soy found in some brands), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before taking this medication, tell your doctor or pharmacist your medical history, especially of: iron overload disorder (such as hemochromatosis, hemosiderosis), use/abuse of alcohol, liver problems, stomach/intestinal problems (such as ulcer, colitis). If your brand of multivitamin also contains folic acid, be sure to tell your doctor or pharmacist if you have vitamin B12 deficiency (pernicious anemia) before taking it. Folic acid may affect certain lab tests for vitamin B12 deficiency without treating this anemia. Untreated vitamin B12 deficiency may result in serious nerve problems (such as peripheral neuropathy). Consult your doctor or pharmacist for details. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant before using this medication. This medication passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: stomach pain, nausea, vomiting, diarrhea. If your doctor has prescribed this medication, do not share it with others. Keep all medical and lab appointments. Some brands may also contain ingredients such as docusate. Ask your doctor or pharmacist if you have questions about the ingredients in your brand. This product is not a substitute for a proper diet. Remember that it is best to get your vitamins and minerals from healthy foods. Maintain a well-balanced diet and follow any dietary guidelines as directed by your doctor. If you are taking this product on a prescribed schedule and miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. This product can decrease the absorption of other drugs such as bisphosphonates (for example, alendronate), levodopa, penicillamine, quinolone antibiotics (for example, cipro 500mg, levofloxacin), thyroid medications (for example, levothyroxine), and tetracycline antibiotics (for example, doxycycline, minocycline such as amoxil, zithromax, flagyl). Separate your doses of these medications as far as possible from your doses of this product. Ask your doctor or pharmacist about how long you should wait between doses and for help finding a dosing schedule that will work with all your medications. If your brand of multivitamin also contains folic acid, be sure to tell your doctor or pharmacist if you take certain anti-seizure drugs (such as hydantoins such as phenytoin). This medication may interfere with certain lab tests, possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

Side Effects

Uses

Interactions

Other Details
