

iMedix: Your Personal Health Advisor.

Cetrotide Kit

Search for medical details concerning Cetrotide subcutaneous on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Nausea, headache, or pain/bruising/redness/swelling/itching at the injection site may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication may cause a condition known as ovarian hyperstimulation syndrome (OHSS). This condition may occur during or after treatment. Rarely, serious OHSS causes fluid to suddenly build up in the stomach, chest, and heart area. Get medical help right away if you develop the following side effects: severe pain or swelling in the lower abdominal (pelvic) area, severe nausea/vomiting, sudden/rapid weight gain, or decreased urination. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, such as: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice any other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used by women having certain fertility treatments (controlled ovarian stimulation). Cetrorelix is usually used in combination with other hormones (FSH and hCG). It works by blocking the release of a certain hormone (luteinizing hormone). Cetrorelix stops eggs from being released too early and gives the eggs time to grow properly. How to use Cetrotide Kit Read the Patient Information Leaflet if available from your pharmacist before you start using cetrorelix and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Learn all preparation and usage instructions from your health care professional and the product package. Follow the instructions for adding the mixing liquid to the medication. Gently shake the vial to mix. Once mixed, use the liquid right away. Before using, check this product visually for particles or discoloration. If either is present, do not use the liquid. Inject this medication under the skin as directed by your doctor, usually once a day for several days. It should be injected in the stomach area, at least one inch away from the belly button. Carefully follow your doctor's instructions on when to start and stop this medication. Before injecting each dose, clean the injection site with rubbing alcohol. Change the injection site each time to lessen injury under the skin. Use this medication exactly as directed by your doctor to get the most benefit from it. Do not increase your dose or use this drug more often or for longer than prescribed. Learn how to store and discard medical supplies

safely. Precautions Before using cetrorelix, tell your doctor or pharmacist if you are allergic to it; or to other products containing gonadotropin releasing hormone (GnRH); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: history of severe allergic reactions, kidney disease. Before having surgery, tell your doctor or dentist about all the products you use (such as prescription drugs, nonprescription drugs, and herbal products). Stop using this medication when you become pregnant. This medication must not be used during pregnancy. If you think you may be pregnant, tell your doctor right away. It is unknown if this drug passes into breast milk. Because of the possible risk to the infant, breast-feeding while using this drug is not recommended. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (such as prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Does Cetrotide Kit interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as vaginal ultrasound, hormone levels) may be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Do not double the dose to catch up. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details
