

# iMedix: Your Personal Health Advisor.

## Cipro Tablet

---

Cipro Tablet is a form of the antibiotic ciprofloxacin, which is part of the fluoroquinolone class. It's used to treat a wide range of bacterial infections, including urinary tract infections, respiratory infections, and skin infections.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

## Description

Side Effects Nausea, diarrhea, dizziness, lightheadedness, headache, and trouble sleeping may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: unusual bruising/bleeding, signs of a new infection (such as sore throat that doesn't go away, fever), signs of kidney problems (such as change in the amount of urine, red/pink urine), signs of liver problems (such as nausea/vomiting that doesn't stop, unusual tiredness, stomach/abdominal pain, yellowing eyes/skin, dark urine). Get medical help right away if you have any very serious side effects, including: severe dizziness, fainting, fast/irregular heartbeat, signs of a tear/break in the main blood vessel called the aorta (such as sudden/severe pain in the stomach/chest/back, shortness of breath). This medication may rarely cause a severe intestinal condition due to a bacteria called *C. difficile*. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. Use of this medication for prolonged or repeated periods may result in oral thrush or a new yeast infection. Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge, or other new symptoms. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat a variety of bacterial infections. Cipro belongs to a class of drugs called quinolone antibiotics. It works by stopping the growth of bacteria. This antibiotic treats only bacterial infections. It will not work for virus infections (such as common cold, flu). Using any antibiotic when it is not needed can cause it to not work for future infections. How to use Cipro Read the Medication Guide and, if available, the Patient Information Leaflet provided by your pharmacist before you start taking ciprofloxacin 500 and each time you get a refill. If you have any questions,

ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, usually twice a day in the morning and evening. The tablet may have a bitter taste if you chew or crush it before taking it. Do not split the tablets unless they have a score line and your doctor or pharmacist tells you to do so. The manufacturer recommends swallowing the whole or split tablet without crushing or chewing. The dosage and length of treatment is based on your medical condition and response to treatment. Drink plenty of fluids while taking this medication unless your doctor tells you otherwise. Take this medication at least 2 hours before or 6 hours after taking other products that may bind to it, decreasing its effectiveness. Ask your pharmacist about the other products you take. Some examples include: quinapril, sevelamer, sucralfate, vitamins/minerals (including iron and zinc supplements), and products containing magnesium, aluminum, or calcium (such as antacids, didanosine solution, calcium supplements). Calcium-rich foods, including dairy products (such as milk, yogurt) or calcium-enriched juice, can also decrease the effect of this medication. Take this medication at least 2 hours before or 6 hours after eating calcium-rich foods, unless you are eating these foods as part of a larger meal that contains other (non-calcium-rich) foods. These other foods decrease the calcium binding effect. Ask your doctor or pharmacist about safely using nutritional supplements/replacements with this medication. For the best effect, take this antibiotic at evenly spaced times. To help you remember, take this medication at the same time(s) every day. Continue to take this medication until the full prescribed amount is finished, even if symptoms disappear after a few days. Stopping the medication too early may result in a return of the infection. Tell your doctor if your condition lasts or gets worse. Warnings Quinolone antibiotics (including ciprofloxacin) may cause serious and possibly permanent tendon damage (such as tendonitis, tendon rupture), nerve problems in the arms and legs (peripheral neuropathy), and nervous system problems. Get medical help right away if you have any of the following symptoms: pain/numbness/burning/tingling/weakness in your arms/hands/legs/feet, changes in how you sense touch/pain/temperature/vibration/body position, severe/lasting headache, vision changes, shaking (tremors), seizures, mental/mood changes (such as agitation, anxiety, confusion, hallucinations, depression, rare thoughts of suicide). Tendon damage may occur during or after treatment with this medication. Stop exercising, rest, and get medical help right away if you develop joint/muscle/tendon pain or swelling. Your risk for tendon problems is greater if you are over 60 years of age, if you are taking corticosteroids (such as prednisone), or if you have a kidney, heart, or lung transplant. This medication may make a certain muscle condition (myasthenia gravis) worse. Tell your doctor right away if you have new or worsening muscle weakness (such as drooping eyelids, unsteady walk) or trouble breathing. Discuss the risks and benefits with your doctor before using this medication. Precautions Before taking ciprofloxacin, tell your doctor or pharmacist if you are allergic to it; or to other quinolone antibiotics (such as amoxicil, zithromax, flagyl, levofloxacin, cipro antibiotic); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: diabetes, heart problems (such as recent heart attack), joint/tendon problems (such as tendonitis, bursitis), kidney disease, liver disease, mental/mood disorders (such as depression), myasthenia gravis, nerve problems (such as peripheral neuropathy), seizures, conditions that increase your risk of seizures (such as brain/head injury, brain tumors, cerebral atherosclerosis), blood vessel problems (such as aneurysm or blockage of the aorta or other blood vessels, hardening of the arteries), high blood pressure, certain genetic conditions (Marfan syndrome, Ehlers-Danlos syndrome). Cipro drug may cause a condition that affects the heart rhythm (QT prolongation). QT prolongation can rarely cause serious (rarely fatal) fast/irregular heartbeat and other symptoms (such as severe dizziness, fainting) that need medical attention right away. The risk of QT prolongation may be increased if you have certain medical conditions or are taking other drugs that may cause QT prolongation. Before using ciprofloxacin, tell your doctor or pharmacist of all the drugs you take and if you have any of the following conditions: certain heart problems (heart failure, slow heartbeat, QT prolongation in the EKG), family history of certain heart problems (QT prolongation in the EKG, sudden cardiac death). Low levels of potassium or magnesium in the blood may also increase your risk of QT prolongation. This risk may increase if you use certain drugs (such as diuretics/"water pills") or if you have conditions such as severe sweating, diarrhea, or vomiting. Talk to your doctor about using ciprofloxacin safely. This medication may rarely cause serious changes in blood sugar, especially if you have diabetes. Check your blood sugar regularly as directed and share the results with your doctor. Watch for symptoms of high blood sugar such as increased thirst/urination. Cipro hcl 500 mg may

increase the blood-sugar-lowering effects of the medication glyburide. Also watch for symptoms of low blood sugar such as sudden sweating, shaking, fast heartbeat, hunger, blurred vision, dizziness, or tingling hands/feet. It is a good habit to carry glucose tablets or gel to treat low blood sugar. If you don't have these reliable forms of glucose, rapidly raise your blood sugar by eating a quick source of sugar such as table sugar, honey, or candy, or by drinking fruit juice or non-diet soda. Tell your doctor right away about the reaction and the use of this product. To help prevent low blood sugar, eat meals on a regular schedule, and do not skip meals. Your doctor may need to switch you to another antibiotic or adjust your diabetes medications if any reaction occurs. This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). This medication may make you more sensitive to the sun. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned or have skin blisters/redness. Cipro may cause live bacterial vaccines (such as typhoid vaccine) to not work well. Tell your health care professional that you are using ciprofloxacin before having any immunizations/vaccinations. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Children may be more sensitive to the side effects of this drug, especially joint/tendon problems. Older adults may be at greater risk for tendon problems (especially if they are also taking corticosteroids such as prednisone or hydrocortisone), QT prolongation, and a sudden tear/break in the main blood vessel (aorta). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This medication passes into breast milk and may have undesirable effects on a nursing infant. Consult your doctor before breast-feeding.

**Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to. Lab and/or medical tests (such as kidney function, blood counts, cultures) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. Do not change brands of this medication without checking with your doctor or pharmacist. Not all brands have the same effects. If you miss a dose, take it as soon as you remember. If it is less than 6 hours before the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: "blood thinners" (such as acenocoumarol, warfarin), strontium. Many drugs besides ciprofloxacin may affect the heart rhythm (QT prolongation), including amiodarone, dofetilide, quinidine, procainamide, sotalol, among others. This medication can slow down the removal of other medications from your body, which may affect how they work. Examples of affected drugs include domperidone, duloxetine, fezolinetant, pifenidone, tasimelteon, tizanidine, among others. Avoid drinking large amounts of beverages containing caffeine (coffee, tea, colas), eating large amounts of chocolate, or taking over-the-counter products that contain caffeine. This drug may increase and/or prolong the effects of caffeine.

---

---

## Side Effects

---

## Uses

---

---

---

---

---

---

---

---

## Interactions

---

---

## Other Details

---

---