

# iMedix: Your Personal Health Advisor.

## Clobex Lotion

---

Clobex Lotion is a potent topical medication that contains clobetasol propionate, a corticosteroid used to treat severe skin conditions such as psoriasis and eczema. This lotion is designed for application on the skin to reduce inflammation, redness, and itching.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

## Description

Side Effects Burning, stinging, itching, irritation, dryness, or redness may occur when this medication is first applied to the skin. This should disappear in a few days as your body adjusts to the medication. If any of these effects last or get worse, notify your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: stretch marks (striae), skin thinning/discoloration, excessive hair growth, acne, small red bumps on the skin (folliculitis). Skin infections can become worse when using this medication. Notify your doctor if redness, swelling or irritation does not improve. Rarely, it is possible this medication will be absorbed from the skin into the bloodstream. This can lead to side effects of too much corticosteroid. These side effects are more likely in children, and in people who use this medication for a long time or over large areas of the skin. Tell your doctor right away if any of the following side effects occur: unusual/extreme tiredness, weight loss, headache, swelling ankles/feet, increased thirst/urination, vision problems. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used by adults to treat a variety of skin conditions (such as eczema, dermatitis, psoriasis). Clobetasol reduces the swelling, itching and redness that can occur in these types of conditions. This medication is a very strong (super-high-potency) corticosteroid. This medication is not recommended for use in children younger than 18 years of age due to an increased risk of side effects. Ask the doctor or pharmacist for details. How to use Clobex topical Read the Patient Information Leaflet if available from your pharmacist before you start using clobetasol lotion and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Use this medication on the skin only. However, do not use it on the face, groin or underarms unless directed to do so by your doctor. Wash and dry your hands before applying. Clean and dry the affected area. Apply a thin film of the medication to the affected area and gently rub in as directed by your doctor, usually 2 times daily. Do not

bandage, wrap or cover the area unless directed to do so by your doctor. After applying the medication, wash your hands, unless you are using this medication to treat the hands. When applying this medication near the eyes, avoid getting it in the eyes because this may worsen or cause glaucoma. Also, avoid getting the medication in your nose or mouth. If you get the medication in your eyes, nose, or mouth, rinse with plenty of water. Do not apply more than 50 grams (50 milliliters or 1.75 ounces) in a given week unless directed by your doctor. Doing so increases the risk of serious side effects. Use this medication only for the condition prescribed. Do not use for longer than 4 weeks unless directed by your doctor. If no improvement occurs in 2 weeks of treatment with clobetasol, or if your condition worsens, stop using this medication and inform your doctor. Precautions Before using clobetasol, tell your doctor or pharmacist if you are allergic to it; or to any other steroids (such as hydrocortisone, dexamethasone); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: poor blood circulation, diabetes, other skin conditions (such as rosacea, perioral dermatitis), immune system problems. Do not use if there is an infection or sore in the area to be treated. Rarely, using corticosteroid medications for a long time or over large areas of skin can make it more difficult for your body to respond to physical stress. Before having surgery or emergency treatment, or if you get a serious illness/injury, tell your doctor or dentist that you are using this medication or have used this medication within the past few months. Though it is unlikely, this medication may temporarily slow down a child's growth if used for a long time. See the doctor regularly so your child's height can be checked. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk when applied to the skin. Similar medications pass into breast milk when taken by mouth. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. This medication can be harmful if swallowed. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for other skin problems unless told to do so by your doctor. A different medication may be necessary in those cases. Lab and/or medical tests (such as adrenal gland function) may be done while you are using this medication, especially if you use this drug for an extended period of time or apply it over large areas of the body. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Avoid freezing. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

**Interactions**

---

---

**Other Details**

---

---