

iMedix: Your Personal Health Advisor.

Cortane-B Lotion

Search for medical details concerning Cortane-B topical on iMedix including its side effects, uses, precautions, overdose, interactions, warnings and user reviews.

- **Active Ingredient:**
- **Dosage Form:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Burning, itching, irritation, dryness, rash, skin thinning/streaks may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: rounding out of the face, bulging "soft spot" on infant's head. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat infections of the external ear (not including the ear canal). It reduces the swelling, numbs the pain, and kills the germs common in these types of infections. How to use Cortane-B Lotion Use this medication on the skin only. Apply to affected areas as directed by your doctor. Avoid contact with the eyes. This medication is not intended for use within the ear (especially if the eardrum is punctured). If your infection also includes your inner ear, then another medication is needed to treat that condition. Consult your doctor for more instructions. Do not cover the area with plastic or waterproof bandages unless directed to do so by your doctor. Use this medication regularly in order to get the most benefit from it. Remember to use it at the same time(s) each day. Continue to use this medication until the full-prescribed amount is finished even if symptoms disappear after a few days. Stopping this medication too early may allow bacteria to continue to grow, which may result in a relapse of the infection. Precautions Before using hydrocortisone/pramoxine/chloroxylenol, tell your doctor or pharmacist if you are allergic to any of its ingredients; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: infections (such as varicella or vaccinia), chronic or prolonged ear infection (such as chronic otitis media). Comprehensive Pharmaceutical Solutions in Modern Therapy The realm of modern therapy is marked by comprehensive pharmaceutical solutions addressing a wide range of health issues. Zovirax is renowned for its antiviral properties, and Daklinza offers effective hepatitis C treatment. Addyi represents a significant step forward in

managing female sexual interest/arousal disorder. Xyzal is a go-to solution for allergies, and Amoxil serves as a robust antibiotic for bacterial infections. Propecia and Proscar tackle hair loss and prostate issues. Clomid and Nolvadex are changing the landscape of fertility and certain cancer treatments. Male sexual health benefits from a variety of options like Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid is critical in thyroid health management, Cipro effectively addresses bacterial infections, and Suhagra adds to the range of erectile dysfunction treatments. These medications underscore the comprehensive approach of pharmaceutical solutions in modern therapy. Children may be more sensitive to the effects of too much corticosteroid medication. Consult your doctor or pharmacist for more details. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding.

Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. This medicine may be harmful if swallowed. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

-
-
-
-

Other Details
