

iMedix: Your Personal Health Advisor.

Dimenhydrinate Vial

Search for medical details concerning dimenhydrinate injection on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects

Drowsiness, constipation, blurred vision, or dry mouth/nose/throat may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. To relieve dry mouth, suck on (sugarless) hard candy or ice chips, chew (sugarless) gum, drink water or use a saliva substitute. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: mental/mood changes (such as restlessness, confusion), fast/irregular heartbeat, shaking (tremor), difficulty urinating. Get medical help right away if you have any very serious side effects, including: seizures. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

Uses

Dimenhydrinate is an antihistamine used to prevent and treat nausea, vomiting and dizziness caused by motion sickness when medication cannot be given by mouth. Dimenhydrinate injection should not be used in newborns because of an increased risk of side effects. How to use Dimenhydrinate Vial This medication is injected into a vein (IV), or it can be injected into a muscle by a health care professional, as directed by your

doctor. Your dosage is based on your medical condition and response to therapy. Tell your doctor if your condition lasts or gets worse.

Interactions

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: antihistamines applied to the skin (such as diphenhydramine cream, ointment, spray). Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or other antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely. This medication may interfere with certain lab tests (such as allergy skin test), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug. Does Dimenhydrinate Vial interact with other drugs you are taking? Enter your medication into the iMedix interaction checker

Other Details

Before using dimenhydrinate, tell your doctor or pharmacist if you are allergic to it; or to diphenhydramine; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: breathing problems (such as asthma, emphysema), glaucoma, stomach problems (such as ulcers, obstruction), difficulty urinating (for example, due to an enlarged prostate gland), heart disease, high blood pressure, liver disease, seizures, overactive thyroid. This drug may make you drowsy or blur your vision. Alcohol or marijuana (cannabis) can make you more drowsy. Do not drive, use machinery, or do anything that needs alertness or clear vision until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Children may be more sensitive to the side effects of this drug. This drug can often cause excitement in young children instead of drowsiness. Older adults may be more sensitive to the side effects of this drug, especially drowsiness, confusion, constipation, or trouble urinating. Drowsiness and confusion can increase the risk of falling. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This drug passes into breast milk and may have undesirable effects on a nursing infant. Consult your doctor before breast-feeding. If someone has

overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: severe drowsiness, seizures, widened pupils. In children, mental/mood changes (such as restlessness, irritability, hallucinations) may occur before drowsiness. Do not share this product with others. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Not applicable. This medication is given in a hospital or clinic and will not be stored at home.
