

iMedix: Your Personal Health Advisor.

Eligard 45 Mg (6 Month) Subcutaneous Syringe

Search for medical details concerning Eligard (6 month) subcutaneous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Hot flashes (flushing), increased sweating, night sweats, tiredness, swelling of the ankles/feet, increased urination at night, dizziness, or mild burning/pain/bruising at the injection site may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Rarely, shrinking of the testicles, breast tenderness/swelling, or reduced sexual interest/ability may also occur as a result of lowered testosterone levels. Talk to your doctor if these effects occur. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: new/worsening bone pain, easily broken bones, increased thirst/urination, mental/mood changes (such as depression, thoughts of suicide, mood swings, aggression). Get medical help right away if you have any very serious side effects, including: symptoms of a heart attack (such as chest/jaw/left arm pain, shortness of breath, unusual sweating), signs of a stroke (such as weakness on one side of the body, trouble speaking, sudden vision changes, confusion), seizures, fast/irregular heartbeat, severe dizziness, fainting. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

How to use Eligard 45 Mg (6 Month) Subcutaneous Syringe Read the Patient Information Leaflet if available from your pharmacist before you start using leuprolide and each time you get a refill. If you have any questions, ask your doctor or pharmacist. This medication is given by injection under the skin as directed by your doctor, usually once every 6 months. This product slowly releases the medication into your blood over a 6-month period. If you are using this medication at home, learn all preparation and usage instructions from your health care professional and the product package. If your product has been kept in the refrigerator, take it out of the refrigerator and allow it to come to room temperature. Before using, check this product visually for particles or discoloration. If either is present, do not use the liquid. Before injecting each dose, clean the injection site with rubbing alcohol. Change the injection site each time to lessen injury under the skin. Learn how to store and discard medical supplies safely. Use this medication regularly to get the most benefit from it. To help you remember, mark

your calendar to keep track of when to receive the next dose. During the first few weeks of treatment, your hormone levels will actually go up before they go down. This is a normal response to this medication. Your symptoms may get worse for a few weeks. If you have prostate cancer that has spread to the spine or caused urinary blockage, you may require closer monitoring by your doctor, especially when you first start treatment. Tell your doctor right away if you have any bone pain, numbness/tingling/weakness of the arms/legs, blood in the urine, painful/difficult urination, unusual weakness, or inability to move. Precautions Before using leuprolide, tell your doctor or pharmacist if you are allergic to it; or to similar drugs (such as histrelin, triptorelin); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: diabetes, heart disease (such as heart attack), stroke, high cholesterol, mental/mood problems (such as depression), seizures. Leuprolide may weaken your bones and increase your risk for bone loss (osteoporosis) if used for a long time. Before using this medication, tell your doctor or pharmacist if you have osteoporosis or if you have any of the following risk factors for osteoporosis: long-term alcohol use, smoking, family history of osteoporosis and broken bones, use of certain medications (for example, corticosteroids such as prednisone, certain anti-seizure drugs such as phenytoin). Leuprolide may cause a condition that affects the heart rhythm (QT prolongation). QT prolongation can rarely cause serious (rarely fatal) fast/irregular heartbeat and other symptoms (such as severe dizziness, fainting) that need medical attention right away. The risk of QT prolongation may be increased if you have certain medical conditions or are taking other drugs that may cause QT prolongation. Before using leuprolide, tell your doctor or pharmacist of all the drugs you take and if you have any of the following conditions: certain heart problems (heart failure, slow heartbeat, QT prolongation in the EKG), family history of certain heart problems (QT prolongation in the EKG, sudden cardiac death). Low levels of potassium or magnesium in the blood may also increase your risk of QT prolongation. This risk may increase if you use certain drugs (such as diuretics/"water pills") or if you have conditions such as severe sweating, diarrhea, or vomiting. Talk to your doctor about using leuprolide safely. This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Older adults may be more sensitive to the side effects of this drug, especially QT prolongation (see above). This medication can affect fertility in males. Ask your doctor for more details. This medication must not be used during pregnancy. It may harm an unborn baby. It is important to prevent pregnancy while using this medication. Consult your doctor for more details and to discuss using reliable forms of non-hormonal birth control (such as condoms, diaphragm with spermicide) while using this medication. If you become pregnant or think you may be pregnant, tell your doctor right away. It is unknown if leuprolide passes into breast milk. Because of the possible risk to the infant, breast-feeding while using this drug is not recommended. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. This medication may interfere with certain lab tests, possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as blood testosterone level, PSA blood test, blood glucose) should be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. It is important to get each dose of this medication as scheduled. If a dose is missed, ask the doctor or pharmacist right away for a new dosing schedule. Do not double the dose to catch up. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details
