

# iMedix: Your Personal Health Advisor.

## Ellura

---

Search for medical details concerning Ellura oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

## Description

Side Effects Stomach upset or diarrhea may occur with very large doses. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Tell your doctor right away if you have any serious side effects, including: severe stomach/abdominal pain. A very serious allergic reaction to this product is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Cranberry has been used for reducing the risk of “bladder infections” (urinary tract infections). It has also been used for decreasing the smell of urine in people who are unable to control urination (incontinent). This product should not be used alone to treat bladder infections. It may not work, and the delay could allow the infection to worsen. Some herbal/diet supplement products have been found to contain possibly harmful impurities/additives. Check with your pharmacist for more details about the brand you use. The FDA has not reviewed this product for safety or effectiveness. Consult your doctor or pharmacist for more details. How to use Ellura Take this product by mouth as directed. Follow all directions and warnings on the product package. If you have any questions, ask your doctor or pharmacist. If you are taking a chewable form of this product, chew thoroughly before swallowing. Tell your doctor if your condition lasts or gets worse. Also, tell your doctor if you have any symptoms of a urinary tract/bladder infection, including fever or burning/painful/frequent urination. You may need medication (such as amoxil, zithromax, cipro, flagyl, levofloxacin) to treat your infection. If you think you may have a serious medical problem, get medical help right away. Precautions Before taking cranberry, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. If you have any of the following health problems, consult your doctor or pharmacist before using this product: current/past kidney stones. Liquid products and chewable forms of this product may contain sugar. Liquid forms may also contain alcohol. Caution is advised if you have diabetes, liver disease, or any other condition that requires you to limit/avoid these substances in your diet. Ask your doctor or pharmacist about using this product safely. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs,

nonprescription drugs, and herbal products). During pregnancy, this product should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this product passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor’s approval. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Keep all medical and lab appointments. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Refer to storage information printed on the package. If you have any questions about storage, ask your pharmacist. Keep all medications and herbal products away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

**Side Effects**

---

---

**Uses**

---

---

---

---

---

---

---

---

**Interactions**

---

---

**Other Details**

---

---