

iMedix: Your Personal Health Advisor.

Enskyce

Search for medical details concerning Enskyce oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Nausea, vomiting, headache, bloating, breast tenderness, swelling of the ankles/feet (fluid retention), or weight change may occur. Vaginal bleeding between periods (spotting) or missed/irregular periods may occur, especially during the first few months of use. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If you miss 2 periods in a row (or 1 period if the pill has not been used properly), contact your doctor for a pregnancy test. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication may raise your blood pressure. Check your blood pressure regularly and tell your doctor if the results are high. Tell your doctor right away if you have any serious side effects, including: lumps in the breast, mental/mood changes (such as new/worsening depression), severe stomach/abdominal pain, unusual changes in vaginal bleeding (such as continuous spotting, sudden heavy bleeding, missed periods), dark urine, yellowing eyes/skin. This medication may rarely cause serious (sometimes fatal) problems from blood clots (such as deep vein thrombosis, heart attack, pulmonary embolism, stroke). Get medical help right away if any of these side effects occur: chest/jaw/left arm pain, confusion, sudden dizziness/fainting, pain/swelling/warmth in the groin/calf, trouble speaking, sudden shortness of breath/rapid breathing, unusual headaches (including headaches with vision changes/lack of coordination, worsening of migraines, sudden/very severe headaches), unusual sweating, weakness on one side of the body, vision problems/changes (such as double vision, partial/complete blindness). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. What is Enskyce? Enskyce is an oral contraceptive pill used to prevent pregnancy. It contains a combination of female hormones (ethinyl estradiol and desogestrel) that prevent ovulation. How effective is Enskyce in preventing pregnancy? When taken correctly, Enskyce is over 99% effective in preventing pregnancy. However, typical use, which accounts for human error, may reduce effectiveness. How should Enskyce be taken? Enskyce should be taken as directed by a healthcare provider. It's usually taken once daily, at the same time every day, for 28 days, followed by a new pack without interruption. What are the possible side effects of Enskyce? Common side effects include nausea, vomiting, headache, bloating, breast tenderness, and changes in menstrual cycle. Serious side effects can include blood clots, stroke, and heart attack, although these are rare. Can Enskyce be used as emergency contraception? Enskyce is not intended for use as emergency contraception. For emergency contraception, other specific

medications are recommended. Can you smoke while taking Enskyce? Smoking increases the risk of serious cardiovascular side effects from hormonal birth control, especially if you are over 35. It is strongly advised not to smoke while taking Enskyce. What should I do if I miss a dose of Enskyce? If you miss a dose of Enskyce, follow the instructions provided in the package insert. Generally, you should take the missed pill as soon as you remember. Missing a pill increases the risk of pregnancy. Can Enskyce help with menstrual cramps or acne? Yes, Enskyce can help in reducing menstrual cramps and treating acne in some women, in addition to providing birth control. Is it safe to take Enskyce during breastfeeding? Enskyce may affect milk production and composition. It's generally recommended to use non-hormonal methods of birth control or a progestin-only pill during breastfeeding. Always consult with a healthcare provider before starting any medication while breastfeeding. Does Enskyce protect against sexually transmitted infections (STIs)? No, Enskyce does not protect against STIs. Use of condoms is recommended to protect against STIs, even while taking Enskyce. Can antibiotics interfere with the effectiveness of Enskyce? Some antibiotics may reduce the effectiveness of Enskyce. Discuss with your healthcare provider if you are prescribed antibiotics to ensure continued contraception coverage. What should I do if I experience severe side effects while taking Enskyce? If you experience severe side effects like chest pain, severe headache, vision changes, or leg pain, seek immediate medical attention as these may be signs of serious conditions. How does Enskyce differ from other birth control pills? Enskyce may have different types or amounts of hormones compared to other birth control pills. Its effectiveness and side effect profile might differ slightly as a result. Can Enskyce cause weight gain? Weight gain can be a side effect of hormonal birth control pills, including Enskyce, though this varies from person to person. Is it necessary to take a break from Enskyce after long-term use? There is no medical necessity to take a break from using Enskyce unless advised by your healthcare provider. Continuous use is effective and safe for most women. Can I start taking Enskyce at any time? It's best to start taking Enskyce on the first day of your menstrual period or as directed by your healthcare provider. Starting at other times might require a backup method of birth control for a short period. How long does it take for Enskyce to become effective? Enskyce is effective at preventing pregnancy almost immediately if started on the first day of your menstrual cycle. If started at other times, it's usually recommended to use an additional birth control method for the first seven days. What happens if I stop taking Enskyce? If you stop taking Enskyce, you can become pregnant. Your menstrual cycle may also be irregular for a short time. Consult your healthcare provider before stopping any prescribed medication. Are there any restrictions on food or beverages while taking Enskyce? There are no specific food or beverage restrictions while taking Enskyce, but a balanced diet can help manage some side effects like weight gain and bloating. Can Enskyce cause mood changes or depression? Some women may experience mood changes or symptoms of depression while taking hormonal contraceptives like Enskyce. If you experience significant mood changes, consult with your healthcare provider. Can I switch from another birth control pill to Enskyce? Yes, you can switch from another birth control pill to Enskyce. It's best to do this under the guidance of a healthcare provider to ensure continuous contraceptive protection.

Uses This combination hormone medication is used to prevent pregnancy. It contains 2 hormones: a progestin and an estrogen. Besides preventing pregnancy, birth control pills may make your periods more regular, decrease blood loss and painful periods, decrease your risk of ovarian cysts, and also treat acne. Using this medication does not protect you or your partner against sexually transmitted diseases (such as HIV, gonorrhea, chlamydia).

How to use Enskyce Read the Patient Information Leaflet provided by your pharmacist before you start using this product and each time you get a refill. The leaflet contains very important information on when to take your pills and what to do if you miss a dose. If you have any questions, ask your doctor or pharmacist.

Transformative Impact of Modern Medications Modern medications have transformed healthcare, with drugs like Zovirax targeting viral infections and Daklinza combating hepatitis C. Addyi represents a breakthrough in the treatment of female sexual interest/arousal disorder. For allergy sufferers, Xyzal offers effective relief, while Amoxil serves as a key antibiotic for a range of bacterial infections. Hair loss and prostate health are addressed by Propecia and Proscar. Clomid and Nolvadex are revolutionizing fertility treatments and cancer therapies. Male sexual health sees advancements with Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid plays a crucial role in managing thyroid disorders, Cipro is a broad-spectrum antibiotic, and Suhagra expands the treatment options for erectile dysfunction. This array of medications underscores the transformative impact of modern pharmaceuticals in enhancing patient care across various health disciplines. Take this medication by mouth as directed by your doctor, usually once daily. Pick a time of day that is easy for you to remember, and take

your pill at the same time each day. If you are taking the chewable tablet, you may either swallow it whole or chew it thoroughly and swallow. Carefully follow the manufacturer's directions for your brand. It is very important to continue taking this medication exactly as prescribed by your doctor. With certain brands of birth control pills, the amount of estrogen and progestin in each active tablet will vary at different times in the cycle. It is very important that you follow the package instructions to find the first tablet, start with the first tablet in the pack, and take them in the correct order. Do not skip any doses. Pregnancy is more likely if you miss pills, start a new pack late, or take your pill at a different time of the day than usual. Vomiting or diarrhea can prevent your birth control pills from working well. If you have vomiting or diarrhea, you may need to use a back-up birth control method (such as condoms, spermicide). Follow the directions in the Patient Information Leaflet and check with your doctor or pharmacist for more details. Taking this medication after your evening meal or at bedtime may help if you have stomach upset or nausea with the medication. You may choose to take this medication at another time of day that is easier for you to remember. No matter what dosing schedule you use, it is very important that you take this medication at the same time each day, 24 hours apart. Ask your doctor or pharmacist if you have any questions. Your pill pack contains 21 pills with active medication. It may also contain 7 reminder pills with no medication. Take one active pill (with hormones) once daily for 21 days in a row. If you are using a product with 28 tablets, take an inactive pill once daily for 7 days in a row after you have taken the last active pill unless otherwise directed by your doctor. If you are using a product with 21 tablets, do not take any tablets for 7 days unless otherwise directed by your doctor. You should have your period during the fourth week of the cycle. After you have taken the last inactive tablet in the pack or gone 7 days without taking an active tablet, start a new pack the next day whether or not you have your period. If you do not get your period, consult your doctor. If this is the first time you are using this medication and you are not switching from another form of hormonal birth control (such as patch, other birth control pills), take the first tablet in the pack on the first Sunday following the beginning of your menstrual period or on the first day of your period. If your period begins on a Sunday, begin taking this medication on that day. For the first cycle of use only, use an additional form of non-hormonal birth control (such as condoms, spermicide) for the first 7 days to prevent pregnancy until the medication has enough time to work. If you start on the first day of your period, you do not need to use back-up birth control the first week. Ask your doctor or pharmacist about how to switch from other forms of hormonal birth control (such as patch, other birth control pills) to this product. If any information is unclear, consult the Patient Information Leaflet or your doctor or pharmacist. **Warnings** Do not use this medication if you smoke cigarettes/use tobacco and are over 35 years old. Smoking raises your risk of stroke, heart attack, blood clots, and high blood pressure from hormonal birth control (such as the pill, patch, ring). The risk of these serious problems increases with age and with the number of cigarettes you smoke. Do not smoke or use tobacco. **Precautions** Before using this medication, tell your doctor or pharmacist if you are allergic to any estrogens (such as ethinyl estradiol, mestranol) or any progestins (such as norethindrone, desogestrel); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: blood clots (for example, in the legs, eyes, lungs), blood clotting disorders (such as protein C or protein S deficiency), high blood pressure, abnormal breast exam, cancer (especially endometrial or breast cancer), high cholesterol or triglyceride (blood fat) levels, depression, diabetes, family or personal history of a certain swelling disorder (angioedema), gallbladder problems, severe headaches/migraines, heart problems (such as heart valve disease, irregular heartbeat, previous heart attack), history of yellowing eyes/skin (jaundice) during pregnancy or while using hormonal birth control (such as pills, patch), kidney disease, liver disease (including tumors), obesity, stroke, swelling (edema), thyroid problems, unexplained vaginal bleeding. If you have diabetes, this medication may affect your blood sugar. Check your blood sugar regularly as directed and share the results with your doctor. Tell your doctor right away if you have symptoms of high blood sugar such as increased thirst/urination. Your doctor may need to adjust your diabetes medication, exercise program, or diet. Chewable tablets may contain sugar and/or aspartame. Caution is advised if you have diabetes, phenylketonuria (PKU), or any other condition that requires you to limit/avoid these substances in your diet. Tell your doctor if you just had or will be having surgery or if you will be confined to a bed or chair for a long time (such as a long plane flight). These conditions increase your risk of getting blood clots, especially if you are using hormonal birth control. You may need to stop this medication for a time or take special precautions. Before having surgery,

tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication may cause blotchy, dark areas on your face and skin (melasma). Sunlight may worsen this effect. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. If you are nearsighted or wear contact lenses, you may develop vision problems or trouble wearing your contact lenses. Contact your eye doctor if these problems occur. It may take longer for you to become pregnant after you stop taking birth control pills. Consult your doctor. This medication should not be used during pregnancy. If you become pregnant or think you may be pregnant, tell your doctor right away. If you have just given birth or had a pregnancy loss/abortion after the first 3 months, talk with your doctor about reliable forms of birth control, and find out when it is safe to start using birth control that contains a form of estrogen, such as this medication. This medication may decrease breast milk production. A small amount passes into breast milk and may have undesirable effects on a nursing infant. Consult your doctor before breast-feeding.

Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include severe nausea and vomiting, sudden/unusual vaginal bleeding. Do not share this medication with others. Regular complete physical exams which include lab and/or medical tests (such as blood pressure, breast exam, pelvic exam, Pap smear) should be done while you are taking this medication. Follow your doctor's instructions for examining your breasts, and report any lumps right away. Keep all medical and lab appointments. Consult your doctor for more details. Refer to the product package information for advice on missed doses. You may need to use back-up birth control (such as condoms, spermicide) to prevent pregnancy. Ask your doctor or pharmacist if you have any questions. If you often forget to take your pills as directed, contact your doctor to discuss switching to another form of birth control. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: aromatase inhibitors (such as anastrozole, exemestane), fezolinetant, ospemifene, tamoxifen, tizanidine, tranexamic acid, certain combination products used to treat chronic hepatitis C (ombitasvir/paritaprevir/ritonavir with or without dasabuvir). Some drugs may cause hormonal birth control to work less well by decreasing the amount of birth control hormones in your body. This effect can result in pregnancy. Examples include griseofulvin, modafinil, rifamycins (such as rifampin, rifabutin), ritonavir, St. John's wort, drugs used to treat seizures (such as barbiturates, carbamazepine, felbamate, phenytoin, primidone, topiramate), HIV drugs (such as nelfinavir, nevirapine), among others. Tell your doctor when you start any new drug, and discuss if you should use additional reliable birth control. Also tell your doctor if you have any new spotting or breakthrough bleeding, because these may be signs that your birth control is not working well. This medication may interfere with certain lab tests (such as blood clotting factors, thyroid), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

Side Effects

Uses

Interactions

-
-
-
-

Other Details
