

iMedix: Your Personal Health Advisor.

Enzymatic Digestant Tablet

Search for medical details concerning Enzymatic Digestant oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects

Diarrhea, abdominal pain/cramps, or nausea may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: severe abdominal discomfort, frequent/painful urination, joint pain. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

Uses

This medication contains digestive enzymes, which are natural substances needed by the body to help break down and digest food. It is used when the pancreas cannot make or does not release enough digestive enzymes into the gut to digest the food. Depending on the amount of enzymes in your product, it may be used for indigestion, as a supplement, or as replacement therapy (such as in chronic pancreatitis, cystic fibrosis, cancer of the pancreas, after surgery on the pancreas or gut). Some supplement products have been found to contain possibly harmful impurities/additives. Check with your pharmacist for more details about

the brand you use. The FDA has not reviewed this product for safety or effectiveness. Consult your doctor or pharmacist for more details. **How to use Enzymatic Digestant Tablet** Take this medication by mouth with meals and snacks as directed by your doctor. The dosage is based on your medical condition, diet, and response to treatment. If you are using the tablet form of the medication, make sure not to keep it in the mouth because doing so can cause irritation to the gums and cheeks. Swallow the medication with water. For the chewable tablets, chew thoroughly before swallowing. If you are using the capsule form of the medication and swallowing is difficult, the capsule may be opened and the powder mixed with food or liquid. Be careful not to inhale any of the powder because doing so can irritate the inside of the nose or cause an asthma attack. Use this medication regularly to get the most benefit from it. If your doctor has recommended that you follow a special diet, it is very important to follow the diet to get the most benefit from this medication. Do not change brands or dosage forms of this product without consulting your doctor or pharmacist. Different products may contain different amounts of the digestive enzymes. Tell your doctor if your condition lasts or gets worse.

Interactions

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: acarbose, miglitol. Does Enzymatic Digestant Tablet interact with other drugs you are taking? Enter your medication into the iMedix interaction checker

Other Details

Before taking this digestive enzyme, tell your doctor or pharmacist if you are allergic to it; or to pork protein; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: sudden/severe swelling of the pancreas (acute pancreatitis), sudden worsening of long-term disease of the pancreas. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this product with others. If you miss a dose, take the next dose with your next meal or snack as directed. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the

bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.
