

# iMedix: Your Personal Health Advisor.

## Ergoloid MESYLATES

---

Search for medical details concerning ergoloid oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Nausea, vomiting, blurred vision, dizziness, drowsiness, flushing, loss of appetite, or stomach upset may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: fainting, slow heartbeat. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat mild problems with mood, memory, concentration, or behavior in older adults. It is believed to increase blood flow to the brain. It is a combination of 3 ergot alkaloids. How to use Ergoloid MESYLATES Take this medication by mouth as directed by your doctor, usually 3 times daily. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same times each day. It may take several weeks before you see any improvement from the medication. Tell your doctor if your condition lasts or gets worse. Precautions Before taking ergoloid mesylates, tell your doctor or pharmacist if you are allergic to them; or to other ergot alkaloids (such as dihydroergotamine, ergotamine); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: mental/mood disorders (such as psychosis), blood pressure problems, heart problems, liver disease. This drug may make you dizzy or drowsy or blur your vision. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness or clear vision until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your

medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Other medications can affect the removal of ergoloid mesylates from your body, which may affect how ergoloid mesylates works. Examples include azole antifungals (such as itraconazole, ketoconazole, voriconazole), cobicistat, macrolide antibiotics (such as amoxil, zithromax, flagyl, levofloxacin, ciprofloxacin), nefazodone, HIV protease inhibitors (such as indinavir, nelfinavir, saquinavir), ritonavir, telithromycin, among others. If you also take "triptan" migraine drugs (such as sumatriptan, rizatriptan), you will need to separate your "triptan" dose from your dose of this medication to reduce the risk of serious side effects. Ask your doctor how long you should wait between your doses of these drugs. Some products have ingredients that could raise your heart rate or blood pressure. Tell your pharmacist what products you are using, and ask how to use them safely (especially cough-and-cold products, diet aids, or other migraine medications). Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Keep all medical appointments. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

