

# iMedix: Your Personal Health Advisor.

## Ethyol Vial

---

Search for medical details concerning Ethyol intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

## Description

**Side Effects** Flushing, chills, unusual feelings of warmth/coldness, blurred vision, dizziness, drowsiness, or pain/redness/swelling at the injection site may occur. Nausea and vomiting occur often and can be severe. Your doctor may prescribe medication to prevent or relieve nausea and vomiting. Eating several small meals, not eating before treatment, or limiting activity may help lessen some of these effects. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: fainting, fast/slow/irregular heartbeat, muscle spasms, mental/mood changes, seizure. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** Amifostine is used to decrease the risk of kidney problems caused by treatment with a certain anti-cancer drug (cisplatin). It is also used to help prevent a certain side effect (dry mouth) caused by radiation treatment for head and neck cancer. Amifostine is thought to work by preventing certain chemicals from damaging healthy cells. It may also help repair cells damaged by cisplatin or radiation treatment. **How to use Ethyol Vial** This medication is given by injection into a vein by a health care professional as directed by your doctor, usually 15 to 30 minutes before cancer treatment. The dosage is based on your medical condition, body size, and response to treatment. To decrease the risk of side effects, drink plenty of fluids while using this medication unless otherwise directed by your doctor. Your doctor may direct you to lie down while receiving this medication to decrease the chances of low blood pressure. Also, if you are taking blood pressure medication, your doctor may direct you to stop taking it for 1 day before receiving this drug. Be sure to ask your doctor when to restart blood pressure medication. **Precautions** Before receiving amifostine, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: low/high blood pressure,

dehydration, heart disease (such as irregular heartbeat, heart failure), kidney disease, previous stroke. This drug may make you dizzy or drowsy or blur your vision. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness or clear vision until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant or plan to become pregnant. Since amifostine is used with anti-cancer treatments which may harm an unborn baby, you should not become pregnant during treatment. Ask about reliable forms of birth control while using this medication with anti-cancer treatments. If you become pregnant, talk to your doctor right away about the risks and benefits of amifostine and the anti-cancer treatments. It is unknown if this medication passes into breast milk. Because of possible risk to the infant, breast-feeding while using this drug is not recommended. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely. Does Ethyol Vial interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Lab and/or medical tests (such as blood pressure, serum calcium levels) should be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Not applicable. This medication is given in a hospital or clinic and will not be stored at home.

## Side Effects

## Uses

## Interactions

---

---

## Other Details

---

---