

# iMedix: Your Personal Health Advisor.

## Evolocumab Wearable Injector

---

Search for medical details concerning evolocumab subcutaneous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Redness, pain, or bruising at the injection site may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How to use Evolocumab Wearable Injector Read the Patient Information Leaflet if available from your pharmacist before you start using evolocumab and each time you get a refill. Carefully read and learn the Instructions for Use. If you have any questions, ask your doctor or pharmacist. Before use, take evolocumab out of the refrigerator. If you are using the prefilled syringe/pen, let it warm up to room temperature for at least 30 minutes. If you are using the on-body infusor, let it warm up to room temperature for at least 45 minutes. Do not heat this medication any other way, such as by heating in the microwave or placing in hot water. Do not shake this medication. Before using, check this product visually for particles or discoloration. If either is present, do not use the liquid. Inject this medication under the skin of the thigh, abdomen, or upper arm as directed by your doctor, usually once every 2 weeks, or once every month. The dosage is based on your medical condition and response to treatment. If your doctor has prescribed the monthly dose and you are using the prefilled syringe/pen, you will use 3 separate injections within 30 minutes. If you are using the on-body infusor, the medication is injected over 5 minutes. Before injecting each dose, clean the injection site with rubbing alcohol. Change the injection site each time to lessen injury under the skin. Do not inject into any areas of skin that are injured, tender, hard, red, or has scars or stretch marks. Use this medication regularly to get the most benefit from it. To help you remember, mark the days on the calendar when you need to receive the medication. Learn how to store and discard medical supplies safely. Precautions Before using evolocumab, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients (such as latex or dry natural rubber found in the needle cover on the prefilled syringe/pen) which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Does Evolocumab Wearable Injector interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as LDL cholesterol levels) should be done while you are using this medication. Keep all medical and lab appointments. If you are using this medication every 2 weeks and miss a dose, use it as soon as you remember if it is within 7 days from the missed dose. If it is longer than 7 days from the missed dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. If you are using this medication once a month and miss a dose, use it as soon as you remember if it is within 7 days from the missed dose, then use your next dose at the regular time. If it is longer than 7 days from the missed dose, use it as soon as you remember and start a new monthly schedule based on this date. Do not double the dose to catch up. Store in the original package in the refrigerator away from light and moisture. Do not freeze. If you keep this medication at room temperature, make sure to use it within 30 days and keep it in the original package. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

---

## **Side Effects**

---

---

---

## **Uses**

---

---

---

---

---

## **Interactions**

---

---

## Other Details

---

---