

# iMedix: Your Personal Health Advisor.

## Evoxac

---

Search for medical details concerning Evoxac oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

## Description

Side Effects Sweating, nausea, runny nose, flushing, frequent urge to urinate, dizziness, weakness, diarrhea, and blurred vision may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. This medication may cause an increase in tears. This can be helpful if you have dry eyes. Tell your doctor if runny eyes become a problem. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: slow/fast/irregular heartbeat, shakiness (tremor), fainting, lung problems (such as increased wheezing/cough/phlegm), mental/mood changes (such as confusion, agitation), severe stomach/abdominal pain. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat symptoms of dry mouth due to a certain immune disease (Sjogren's syndrome). Cevimeline belongs to a class of drugs known as cholinergic agonists. It works by stimulating certain nerves to increase the amount of saliva you produce, making it easier and more comfortable to speak and swallow. How to use Evoxac Take this medication by mouth with or without food as directed by your doctor, usually 3 times daily. Dosage is based on your medical condition and response to treatment. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same times each day. You may continue to drink water or use saliva substitutes as needed for moisture in your mouth. You may start to feel some benefit in 1 to 2 weeks. However, it may take up to 6 weeks to feel the full benefit. Tell your doctor if your condition lasts or gets worse. Precautions Before taking cevimeline, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: breathing problems (such as asthma, chronic bronchitis, chronic obstructive pulmonary disease-COPD), certain eye conditions (such as night blindness, acute iritis, narrow-angle glaucoma), heart disease (such as chest pain, heart failure, heart attack, slow heartbeat), low or

high blood pressure, liver problems, gallbladder disease (such as gallstones), kidney stones, mental/mood disorders (such as depression, psychoses, thinking/understanding problems like dementia, Alzheimer's), stomach problems (such as chronic heartburn, ulcer). This drug may make you dizzy or cause vision problems, especially at night. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness or clear vision, especially at night, until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). If cevimeline makes you sweat heavily, drink plenty of fluids so that you do not become dehydrated. If you are unable to drink enough fluids, talk with your doctor right away. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: beta blockers (such as metoprolol, atenolol). Other medications can affect the removal of cevimeline from your body, which may affect how cevimeline works. Examples include cimetidine, amiodarone, ritonavir, fluoxetine, among others. Does Evoxac interact with other drugs you are taking? Enter your medication into the iMedix interaction checker

**Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: extreme sweating, shakiness, slowed/irregular heartbeat, severe breathing problems, confusion, fainting, stomach/abdominal cramping. Do not share this medication with others. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---