

# iMedix: Your Personal Health Advisor.

## Female Cialis (Tadalafil)

---

Generic Female Cialis (Tadalafil) increases the blood flow to the female genital area after taking just one tablet already. Sexual necking from the partner causes more emotions and feelings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

General information The absence of sexual attraction in women is not only about the hormones, but a psychological problem as well. Any hormone imbalance and psychological stress can provoke sexual dysfunction. Sexual interest may fade away in women of any age. It is very important to maintain the libido at a high level because regular female orgasm improves: brain functions; metabolic process in women body. Female Cialis- is a pink- coloured pill, which has a 20 mg of the main active ingredient – Tadalafil. Regular orgasm prevents the development of various gynecological diseases. Female sexual dysfunction can manifest itself with different symptoms. Some women do not have problems with sexual excitement but they can not get an orgasm no matter how they try. Other women do not have a sexual drive at all. Some women experience discomfort caused by the dryness of the vagina during the sexual intercourse. This causes an inconvenience to the both sexual partners. The etiology of sexual dysfunction in women are: Hormone disruption. A hormone called Testosterone is responsible for libido both in men and in women. The insufficiency of this hormone in the human body becomes the cause of the lack of sexual desire. Psychological and emotional problems. Statistically, the number of women suffering from low libido as a result of psychological disorders over the past few years grown up to 50%. Head injuries and diseases of the central nervous system, in which the part of the brain responsible for the attraction. Some neurological diseases and head injuries can cause disruption of the connections in the brain responsible for sex satisfaction. These include a scarcity or excess of blood flow to the tissues, parasitic disease, benign and malignant diseases, hematomas caused by trauma. Drugs administration. Sexual dysfunction can occur as a side effect when a woman takes antifertility agents, drugs for reduction of arterial blood pressure, tranquilizers, neuroleptics and other medications for the treatment of neurological problems. What is Female Cialis? Female Cialis is a medication intended to enhance sexual function in women. It contains tadalafil, which is also used in the treatment of erectile dysfunction in men. It aims to increase sexual arousal and improve the experience of sexual activity in women. How does Female Cialis work? Female Cialis works by increasing blood flow to the genital area, which may help increase sensitivity and sexual arousal in women. It is thought to work by relaxing smooth muscles and improving blood flow, similar to its mechanism in treating erectile dysfunction in men. What are the benefits of taking Female Cialis? The benefits of taking Female Cialis may include increased sexual desire, enhanced natural lubrication, improved sensitivity in the genital area, and potentially more satisfying sexual experiences. What is the recommended dosage of Female Cialis? The recommended dosage of Female Cialis varies, and it is important to follow a healthcare

provider's prescription. Typically, it is taken before anticipated sexual activity but not more than once daily. Are there any side effects of Female Cialis? Side effects of Female Cialis may include nausea, headaches, dizziness, stomach upset, flushing, and nasal congestion. In rare cases, more severe side effects such as allergic reactions or changes in vision and hearing can occur. Is Female Cialis safe for all women? Female Cialis may not be safe for all women, especially those with certain health conditions like heart disease, kidney or liver problems, or those who are pregnant or breastfeeding. It's crucial to consult a healthcare provider before starting this medication. Can Female Cialis be used with alcohol? It's advisable to limit or avoid alcohol while taking Female Cialis, as alcohol can increase the risk of side effects such as dizziness and can affect sexual functioning. How long does it take for Female Cialis to work? The onset of action for Female Cialis can vary, but it generally takes about 30 minutes to an hour. Its effects can last for up to 36 hours. Does Female Cialis require a prescription? Yes, Female Cialis is a prescription medication, and it's important to consult a healthcare provider to determine if it is appropriate for your condition and to receive a prescription. Can Female Cialis be taken with other medications? Female Cialis can interact with certain medications, especially those used to treat heart conditions and blood pressure. It's essential to inform your healthcare provider about all medications you're taking to ensure safety and prevent harmful interactions.

Female Cialis is a medicine, which helps to solve the problem of female sexual dysfunction. This is a medical product that increases the blood flow to the female genitalia and thus enhances their sensitivity to stimulation. The medicine is made by an Indian pharmaceutical company. The design of Female Cialis is for sale at the market after receiving positive results from the administration of Cialis for men. Which diseases does this medicine help to cure? There are some physiological pathologies in women not allowing to enjoy the sexual intimacy with the partner: The postmenopausal period and the onset of menopause in most cases give the woman unpleasant sensations and reduce her sexual desire, the sensitivity of the genitals and general relaxation during intimacy with a man. Hormonal changes during pregnancy and after childbirth sometimes reduce a partner's sensitivity or sexual intimacy, do not allow to have any pleasure and relaxation. There is a reduction in lubrication of the vagina, dryness of the vagina or bad smell. These signs report about pathology in the woman's reproductive system and, as a result, about a decrease in sexual desire. There is an interference with blood flow in pelvic organs due to injuries and physiological changes. Inadequate blood supply to appendages, vagina, adrenal glands reduces their sensitivity and the corresponding reaction of the reproductive system. There is a psychological fear of sexual intimacy. There is a problem with the production of Testosterone or Estrogen that are basically very important for proper sexual relations of a woman with a man. The active substance of the medicine is Tadalafil (same as in Cialis for men). The drug is available in the form of tablets with different dosages (10, 20 and 30 mg). How to use? Doctors recommend to use the drug for 15 – 30 minutes before sexual activity. This time is enough for the drug to be completely absorbed into the blood. Do not use more than one tablet (20 mg) per day. Action time of the medicine is more than 24 hours. Firstly, the medicine should be taken with water. The effect does not improve with taking two or more tablets instead but there is a risk of side effects. Secondly, dosage can be reduced down to 10 or 5 mg if there are no serious disorders at the level of hormones that regulate the sexual system of women. Thirdly, the substance is absorbed more slowly if a woman drinks alcohol but the effectiveness of the medicine does not change. What effects does the medicine have on the woman's body? Generic Female Cialis (Tadalafil) increases the blood flow to the female genital area after taking just one tablet already. Sexual necking from the partner causes more emotions and feelings. The walls of the vagina become more sensitive, the production of natural lubricant during the sexual encounter starts. This gives a woman the maximum pleasure and high feelings from sex. Tadalafil affects the erogenous zones of women and increases their sensitivity. This makes the sexual act more intense and the orgasm stronger. The medicine starts its action only if there is a sexual attraction to a partner. The medicine causes the following effects: Increased threshold of sensitivity of external genital organs at tactile contact. The effect of the medicine is observed even at the first intake. It helps to liberate oneself and completely surrender to the senses. Full relaxation improves the quality of sexual intimacy, accelerates the onset of orgasm and improves its quality. A woman can feel new emotions. The fact of a failure during sexual intimacy is excluded. Using a Female Cialis during pregnancy The medicine is contraindicated in pregnancy and lactation. Limitations of the medicine administration and side effects The medicine is contraindicated in the following cases: the patient's age is less than 18 years; individual intolerance to Tadalafil; diseases of the blood and cardiovascular system; cardiac decompensation and other severe cardiac pathologies (eg, prior stroke). It is necessary to pass the

ECG and consult a cardiologist before taking the medicine; peptic ulcer; oncological diseases; pregnancy and lactation; simultaneous therapy of other diseases with the use of drugs, which include organic nitrates. When you visit a gynecologist you need to tell in detail about taking other medications; alcohol dependence and even individual approaches to drinking alcohol-containing beverages during treatment with Cialis; prolonged loss of blood. Female Cialis Side effects occur in rare cases. Possible medicine-related adverse events are listed below: headache that can appear because of increased blood flow. This is due to the fact that the active substance dilates the blood vessels. It is necessary to get a consultation of a specialist and decide whether to change the dose of the medicine or choose an analogue if there is a headache with repeated use of the medicine; spasms in the regions of vertebral column, most often in the lumbar part of spine. These side effects usually disappear after 2-3 days of therapy. This side effect arises because the circulation in the pelvic organs increases; unpleasant sensations in the area of the gastrointestinal tract (flatulence, belching, spasm); unpleasant sensations in the nasopharynx (stuffiness in nose, copious mucus secretions); muscular aches; hypersensitivity reaction on the skin (if there is an individual intolerance to the medicine components). Doctors recommend to stop using this medicine and pay attention to other medicines or biological supplements if the side effects do not stop. It is necessary as soon as possible to visit a doctor if there is an increased heart rate, blood pressure changes, a strong allergic reaction accompanied by puffiness. Advantages of the medicinal product Female Cialis is an effective modern medical product that helps to transform intimacy from routine exercise into a real pleasure, pleasant for both partners. The advantages of the drug are listed below: safe and carefully tested composition of a medicine; quick and long-term effect of the medicine (within 24 hours); effective increase in sexual desire even during menopause and after surgery for removal of alvuses; affordable cost; rare adverse drug events; possibility to use with fatty foods. Female Cialis Reviews of doctors and patients The medicine helps to get intimate life going. Reports of adverse events are rare.

---

---

## Side Effects

---

---

## Uses

---

---

---

---

---

---

---

---

## Interactions

---

-

- 
- 
- 

---

## Other Details

---

---