

iMedix: Your Personal Health Advisor.

FVIII Rec,B-Dom Delet Peg-Aucl Solution, Reconstituted (Recon Soln)

Search for medical details concerning antihemophilic FVIII recom,B-domain del peg-aul intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Flushing of the face, headache, nausea, and fast heartbeat may sometimes occur and can be lessened by giving this medication more slowly. Burning/redness/irritation at the injection site, fever, and chills may also occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing, chest discomfort/tightness. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This medication is used to control and prevent bleeding episodes in people with low levels of factor VIII (hemophilia A). It is also used in these patients before surgery to prevent bleeding. People with low factor VIII levels are at risk for bleeding longer after an injury/surgery and for bleeding inside the body (especially into the joints and muscles). This product contains a man-made form of factor VIII, also called antihemophilic factor. This product is used to temporarily replace the missing factor VIII, a protein (clotting factor) that is normally in the blood, so that the blood can clot and the bleeding can stop. This medication should not be used to treat von Willebrand disease. **How to use** FVIII Rec,B-Dom Delet Peg-Aucl Solution, Reconstituted (Recon Soln) This medication is given by injection into a vein as directed by your doctor. The medication may need to be given more slowly over a longer time depending on your dose and how you respond to it. After first receiving this medication in a hemophilia treatment center or hospital, some patients may be able to give this medication to themselves at home. If your doctor directs you to give this medication at home, read the Patient Information Leaflet available from your pharmacist before you start using this medication and each time you get a refill. Learn all preparation and usage instructions in the product package. Learn how to store and discard needles and medical supplies safely. If you have any questions, ask your health care professional. If the medication and the solution used to mix it were refrigerated, bring both to room temperature before combining. After adding the solution to the powder, gently swirl the mixture to completely dissolve the

powder. Do not shake. Before using this product, check it visually for particles or discoloration. If either is present, do not use the liquid. Use right away according to the product instructions. The dosage is based on your medical condition, age, weight, blood test results, and response to treatment. Follow your doctor's instructions closely. Tell your doctor if your condition does not improve or if it worsens. Precautions Before using this medication, tell your doctor or pharmacist if you are allergic to any antihemophilic factor (factor VIII) products; or if you have any other allergies. Some brands may also contain polyethylene glycol (PEG), or inactive ingredients (such as natural rubber/latex), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: a certain type of hemophilia (congenital hemophilia A with inhibitors). Manufacturers of some brands of this medication recommend that you monitor your heartbeat during treatment. If your heart starts to beat faster, it is recommended that you give this medication more slowly or temporarily stop the infusion until your heart rate returns to normal. Consult your doctor for more details. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Does FVIII Rec,B-Dom Delet Peg-Aucl Solution, Reconstituted (Recon Soln) interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab tests (such as factor VIII levels) should be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. It is important to follow the dosing schedule as directed by your doctor. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Do not double the dose to catch up. Store the medication according to the manufacturer's instructions. Ask your pharmacist if you have any questions. Do not freeze this product. Protect the product from light. After mixing, use the product within the time frame stated in the manufacturer's instructions and throw away any unused portion. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details
