

# iMedix: Your Personal Health Advisor.

## Gas Relief Ultra Strength 180 Mg Capsule

---

Search for medical details concerning Gas Relief Ultra Strength oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

**Side Effects** This product usually has no side effects. If you have any unusual effects, contact your doctor or pharmacist promptly. If your doctor has directed you to use this product, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this product do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This product is used to relieve symptoms of extra gas such as belching, bloating, and feelings of pressure/discomfort in the stomach/gut. Simethicone helps break up gas bubbles in the gut. Check the ingredients on the label even if you have used the product before. The manufacturer may have changed the ingredients. Also, products with similar names may contain different ingredients meant for different purposes. Taking the wrong product could harm you. **How to use Gas Relief Ultra Strength 180 Mg Capsule** Follow all directions on the product package. If your doctor has prescribed this medication, take it as directed. If you have any questions, ask your doctor or pharmacist. Take this product by mouth, usually as needed after meals and at bedtime. The dosage is based on your age, medical condition, and response to treatment. Do not increase your dose or take this product more often than directed. Do not take more of this product than recommended for your age. If you are using the chewable tablets, chew each tablet well and swallow. If your condition lasts or gets worse, or if you think you may have a serious medical problem, get medical help right away. **Precautions** Before taking simethicone, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. If you have any health problems, consult your doctor or pharmacist before using this product. Chewable tablets may contain sugar and/or aspartame. Caution is advised if you have diabetes, phenylketonuria (PKU), or any other condition that requires you to limit/avoid these substances in your diet. Ask your doctor or pharmacist about using this product safely. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant before using this product. Simethicone does not pass into

breast milk and is unlikely to harm a nursing infant. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Simethicone can decrease the absorption of thyroid medications (such as levothyroxine). If you take thyroid medication, take it at least 4 hours before or after products that contain simethicone. Does Gas Relief Ultra Strength 180 Mg Capsule interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Keep all medical and lab appointments. If you have problems with gas, it may help to limit foods known to cause gas and bloating (including beans, cabbage, high-fat foods) or use digestive enzymes. Ask your doctor or a dietitian to help review what you eat. Belching may be caused by swallowing too much air. Eating/drinking quickly, chewing gum, and smoking can cause you to swallow more air. If you are taking this product on a regular schedule and miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

---

## **Side Effects**

---

---

---

## **Uses**

---

---

---

## **Interactions**

---

---

---

## **Other Details**

