

iMedix: Your Personal Health Advisor.

Gavreto

Search for medical details concerning Gavreto on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Tiredness, constipation or diarrhea, muscle/joint pain, fever, or swelling in the hands/legs/face may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. People using this medication may have serious side effects. However, you have been prescribed this drug because your doctor has judged that the benefit to you is greater than the risk of side effects. Careful monitoring by your doctor may decrease your risk. This medication may raise your blood pressure. Check your blood pressure regularly and tell your doctor if the results are high. Your doctor may control your blood pressure with medication.

Tell your doctor right away if you have any serious side effects, including: easy bruising/bleeding, slow wound healing, new or worsening lung problems (such as shortness of breath, cough), signs of liver problems (such as nausea/vomiting that doesn't stop, loss of appetite, stomach/abdominal pain, yellowing eyes/skin, dark urine). Pralsetinib sometimes causes side effects due to the rapid destruction of cancer cells (tumor lysis syndrome). To lower your risk, your doctor may add a medication and tell you to drink plenty of fluids. Tell your doctor right away if you have symptoms such as: low back/side pain (flank pain), signs of kidney problems (such as painful urination, pink/bloody urine, change in the amount of urine), muscle spasms/weakness. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **How to use Gavreto** Read the Patient Information Leaflet if available from your pharmacist before you start using pralsetinib and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth on an empty stomach as directed by your doctor, usually once daily. Do not eat food for at least 2 hours before and at least 1 hour after taking pralsetinib. The dosage is based on your medical condition, response to treatment, and other medications you may be taking. Be sure to tell your doctor and pharmacist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). If you have certain side effects, your doctor may adjust your dose or stop treatment for some time. Do not increase your dose or use this drug more often or for longer than prescribed. Your condition will not improve any faster, and your risk of serious side effects will increase. Take this medication regularly to get the most benefit from it. To help you

remember, take it at the same time each day. If you vomit after taking a dose, do not take another dose. Take your next dose at your regular time the next day. Since this drug can be absorbed through the skin and lungs and may harm an unborn baby, women who are pregnant or who may become pregnant should not handle this medication or breathe the dust from the capsules. Precautions Before taking pralsetinib, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: blood/bleeding problems, high blood pressure, liver problems, lung or breathing problems other than lung cancer. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication may cause wounds to heal slowly or poorly. Your doctor or dentist may tell you to temporarily stop treatment with this medication at least 5 days before surgery. Ask your doctor or dentist for specific instructions about when to stop and when to restart treatment with this medication. Tell your doctor/dentist right away if you have wounds that are not healing well. Since this drug can be absorbed through the skin and lungs and may harm an unborn baby, women who are pregnant or who may become pregnant should not handle this medication or breathe the dust from the tablets. Tell your doctor if you are pregnant or plan to become pregnant. You should not become pregnant while using pralsetinib. Pralsetinib may harm an unborn baby. Your doctor should order a pregnancy test before you start this medication. Women using this medication should ask about reliable forms of non-hormonal birth control during treatment and for 2 weeks after the last dose. Men using this medication should ask about reliable forms of birth control during treatment and for 1 week after the last dose. If you or your partner becomes pregnant, talk to your doctor right away about the risks and benefits of this medication. It is unknown if this drug passes into breast milk. Because of the possible risk to the infant, breast-feeding is not recommended while using this drug and for 1 week after the last dose. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Other medications can affect the removal of pralsetinib from your body, which may affect how pralsetinib works. Examples include azole antifungals (such as itraconazole, ketoconazole), drugs used to treat seizures (such as phenytoin, carbamazepine), St. John's wort, among others. Does Gavreto interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as blood mineral levels, kidney/liver function, complete blood counts) should be done before you start taking this medication and while you are taking it. Keep all medical and lab appointments. If you miss a dose, take it as soon as you remember on the same day. If it is the next day when you remember, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Page 1 of 1

Other Details
