

# iMedix: Your Personal Health Advisor.

## Gelclair Gel In Packet

---

Search for medical details concerning Gelclair oral mucous membrane on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

**Side Effects** Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This medication is used to relieve pain from mouth sores and swelling due to various causes including chemotherapy, radiation treatment, dental surgery/devices, and disease. Reducing pain helps you to eat and drink more comfortably. This medication works by forming a protective coating over the inside of the mouth to cover the irritation. It also keeps the mouth moist which promotes healing. **How to use Gelclair Gel In Packet** Open the packet and pour the gel into a glass. Add 1 tablespoonful (15 milliliters) of water and stir the mixture well. If the liquid is too thick, you may add 1 to 2 tablespoonfuls (15 to 30 milliliters) more water. Rinse your mouth thoroughly with the liquid for at least 1 minute or as long as possible, then gargle and spit out. Use this medication as directed by your doctor, usually 3 times a day. Do not eat or drink for at least 1 hour after use. This medication may be used without mixing with water if necessary. Tell your doctor if your condition does not improve after 7 days of treatment. **Precautions** Before using this medication, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history. Tell your doctor if you are pregnant before using this medication. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. **Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. **Does Gelclair Gel In Packet interact with other drugs you are taking?** Enter your medication into the iMedix interaction checker. **Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call

911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from direct sunlight. Do not refrigerate. Do not use any packets that are torn or damaged. Over time, the gel may become a little thicker or darker in color. However, you may still use it as long as the expiration date has not been reached. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---