

iMedix: Your Personal Health Advisor.

Gemcitabine 2,200 Mg/220 Ml (10 Mg/Ml) In 0.9 % Sod.Chlor IV Piggyback

Search for medical details concerning gemcitabine in 0.9 % sodium chloride intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Nausea, vomiting, diarrhea, mouth sores, drowsiness, muscle aches, and pain/redness/swelling at the injection site may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Temporary hair loss may occur. Normal hair growth should return after treatment has ended. People using this medication may have serious side effects. However, you have been prescribed this drug because your doctor has judged that the benefit to you is greater than the risk of side effects. Careful monitoring by your doctor may decrease your risk. Tell your doctor right away if you have any serious side effects, including: signs of lung problems (such as shortness of breath, cough), dizziness/lightheadedness, fainting, swelling ankles/feet/hands, numb/tingling skin, fast/slow/irregular heartbeat, signs of kidney problems (such as change in the amount of urine, bloody urine), signs of liver disease (such as yellowing eyes/skin, dark urine, stomach/abdominal pain). Get medical help right away if you have any very serious side effects, including: chest/jaw/left arm pain, severe headache, trouble speaking, weakness on one side of the body, vision changes, mental/mood changes (such as confusion), seizures. This medication may decrease bone marrow function, an effect that may lead to a low number of blood cells such as red cells, white cells, and platelets. This effect can cause anemia, decrease your body's ability to fight an infection, or cause easy bruising/bleeding. Tell your doctor right away if you develop any of the following symptoms: unusual tiredness, pale skin, signs of infection (such as sore throat that doesn't go away, fever, chills), easy bruising/bleeding. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Gemcitabine is used to treat certain types of cancer (including biliary tract, breast, lung, ovarian, pancreatic). It is a chemotherapy drug that works by slowing or stopping the growth of cancer cells. How to use Gemcitabine 2,200 Mg/220 Ml (10 Mg/Ml) In 0.9 % Sod.Chlor IV Piggyback This medication is given by injection into a vein by a healthcare professional. It is given over 30 minutes as directed by your doctor. The dosage is based on your medical condition, body size, and response to treatment. Precautions Before using gemcitabine, tell your doctor or pharmacist if you are allergic to it; or if you have any other

allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, kidney disease, radiation treatment, heart problems (such as irregular heartbeat, heart failure). This drug may make you drowsy. Alcohol or marijuana (cannabis) can make you more drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Gemcitabine can make you more likely to get infections or may make current infections worse. Stay away from anyone who has an infection that may easily spread (such as chickenpox, COVID-19, measles, flu). Talk to your doctor if you have been exposed to an infection or for more details. Tell your health care professional that you are using gemcitabine before having any immunizations/vaccinations. Avoid contact with people who have recently received live vaccines (such as flu vaccine inhaled through the nose). To lower the chance of getting cut, bruised, or injured, use caution with sharp objects like razors and nail cutters, and avoid activities such as contact sports. Older female adults are more sensitive to the side effects of this medication, especially the risk of infection and easy bruising/bleeding. Tell your doctor if you are pregnant or plan to become pregnant. You should not become pregnant while using gemcitabine. Gemcitabine may harm an unborn baby. Your doctor should order a pregnancy test before you start this medication. Women using this medication should ask about reliable forms of birth control during treatment and for 6 months after the last dose. Men using this medication should ask about reliable forms of birth control during treatment and for 3 months after the last dose. If you or your partner becomes pregnant, talk to your doctor right away about the risks and benefits of this medication. It is unknown if this medication passes into breast milk. Because of the possible risk to the infant, breast-feeding while using this drug and for 1 week after stopping treatment is not recommended. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Does Gemcitabine 2,200 Mg/220 MI (10 Mg/MI) In 0.9 % Sod.Chlor IV Piggyback interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Lab and/or medical tests (such as complete blood count, liver/kidney function, blood pressure) should be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Not applicable. This medication is given in a clinic and will not be stored at home.

Side Effects

Uses

Interactions

Other Details
