

iMedix: Your Personal Health Advisor.

Generic Cialis Black

Generic Cialis Black is an effective drug which will be extremely useful for men who suffer from sexual disorders of any severity stage - advanced, medium or light. The main substance is Tadalafil. The drug must be taken about half an hour before the sexual intercourse. Eating does not affect drug exposure. The duration of the drug activity is approximately 35 hours. During this time a man can have sex without any problems.

- **ActiveIngredient:** Tadalafil
 - **DosageForm:** Tablets
 - **Dosage:** 800 mg
 - **Indications:** Erectile Dysfunction (ED)
 - **Manufacturer:** Centurion Laboratories Pvt Ltd
 - **Storage:** Store in a cool, dry place away from direct sunlight and moisture. Keep out of reach of children.
 - **Market Price:** 1.18
 - **Drug Status:** Prescription Only
-

Description

What is Cialis Black? Cialis Black is a generic version of Cialis, a medication used to treat erectile dysfunction (ED) in men. The active ingredient in Cialis Black is Tadalafil, which helps enhance blood flow to the penis, making it easier to achieve and maintain an erection. What is the difference between Cialis and Cialis Black? Cialis Black is often marketed as a more potent or longer-lasting version of Cialis. However, it's important to note that there's no official "Cialis Black" produced by the manufacturer. The black pills sold under this name might be counterfeit and could contain unknown ingredients.

General Instructions

How to Take Cialis Black (if it's genuine)? Dosage and Frequency: Typically, one pill is taken about 20 to 60 minutes before sexual activity. Do not exceed one dose per day. Method of Consumption: Take the pill with a glass of water. It can be taken with or without food. Timing: The medication should be taken as needed, prior to sexual activity. It does not need to be taken daily unless advised by a healthcare professional. Important Precautions (for genuine Cialis): Medication Interactions: Cialis Black can interact with certain medications, particularly those containing nitrates (often prescribed for chest pain) and medications used to treat high blood pressure. Consult a Doctor: Always consult a healthcare provider before starting Cialis Black to ensure it is safe for you, especially if you have pre-existing health conditions. Dosage Limitations: Do not exceed the recommended dosage as it can increase the risk of side effects. Follow your doctor's instructions carefully.

Side Effects

Common Side Effects: Headache: You might experience a mild to moderate headache after taking Cialis Black. Flushing: This is a warm, red feeling on your face, neck, or chest. Indigestion: You may feel some

discomfort in your stomach or a burning sensation. **Muscle Aches:** You could have some mild pain or discomfort in your muscles. **Nasal Congestion:** Your nose might feel stuffy or blocked. **Serious Side Effects:** **Chest Pain:** If you feel pain in your chest, especially during or after sex, seek medical help immediately. **Sudden Vision Changes:** If you suddenly lose your vision or experience any vision changes, stop taking the medication and contact a doctor right away. **Sudden Hearing Loss:** If you notice a sudden decrease or loss of hearing, discontinue use and get medical attention. **Prolonged Erections:** An erection lasting more than 4 hours can cause permanent damage to your penis. Seek emergency medical care if this occurs. **Severe Allergic Reactions:** Symptoms like rash, itching, or swelling (especially of the face, tongue, or throat) require immediate medical attention. **Side Effects of Counterfeit Cialis Black:** Counterfeit pills may contain harmful ingredients. This can lead to serious health problems, including poisoning, organ damage, or severe allergic reactions. Always ensure you are purchasing medication from a trusted source to avoid these risks.

Uses

Treating Erectile Dysfunction (for genuine Cialis): Cialis Black helps men achieve and maintain an erection by increasing blood flow to the penis. It is important to note that it does not cure erectile dysfunction but provides temporary relief from its symptoms. The medication works only when you are sexually aroused. **Counterfeit Cialis Black – Avoid Use:** Counterfeit versions of Cialis Black can be dangerous. They may contain unknown ingredients that can cause severe health issues. Avoid using counterfeit medications and only obtain Cialis from reputable sources to ensure safety and effectiveness.

Safety advice

Interactions Alcohol:

- Unsafe
- Consuming alcohol while taking Cialis Black 800 mg can increase the risk of side effects like dizziness and low blood pressure. It's best to avoid alcohol.

Interactions Other Medications:

- Consult your doctor
- Cialis Black 800 mg may interact with other medications. Consult your doctor before taking it alongside other drugs to avoid potential risks.

Special Groups Pregnancy:

- Unsafe
- Cialis Black 800 mg is not intended for use during pregnancy. Women who are pregnant should not use this medication.

Special Groups Breastfeeding:

- Unsafe
- This medication is not recommended for breastfeeding women, as its effects on infants are not well studied.

Special Groups Elderly:

- Safe if prescribed
- Elderly individuals can use Cialis Black 800 mg, but only if prescribed by a doctor, as they may be more sensitive to its effects.

Special Groups Children:

- Unsafe
- Cialis Black 800 mg is not suitable for children and should not be used by anyone under 18 years of age.

Effects on Activities Driving:

- Use with caution
- Cialis Black 800 mg may cause dizziness or vision issues, so be cautious when driving after taking this medication.

Effects on Activities Operating Machinery:

- Use with caution
 - Be careful when operating machinery, as Cialis Black 800 mg may affect your ability to do so safely.
-

Concerns

Counterfeit Cialis Black: Significant Risk: Counterfeit pills pose a major health risk. They may contain harmful substances or incorrect dosages. These unregulated pills can lead to unpredictable and dangerous side effects. **Unknown Ingredients:** The contents of counterfeit Cialis Black are not regulated, meaning they could include dangerous or ineffective ingredients. These unknown substances can cause severe allergic reactions or other health issues. **Severe Side Effects:** Using counterfeit medication can lead to serious health problems, including severe side effects, toxicity, and even death. Without knowing what's inside these pills, the risk of adverse reactions is significantly higher. **Ineffectiveness:** Counterfeit medications might not contain any active ingredients or may contain them in incorrect amounts, leading to ineffectiveness in treating erectile dysfunction. **Potential for Addiction:** Some counterfeit pills might contain substances that can cause dependency or other long-term health issues. **Impact on Overall Health:** Using counterfeit drugs can lead to overall health deterioration due to the potential presence of toxic substances. **Reputable Sources:** Always ensure that you are purchasing medications from trustworthy and reputable sources, such as licensed pharmacies. This ensures the medication is safe, effective, and has been tested for quality. **Verify the authenticity of the medication through packaging, holograms, and official approval seals.** **Consulting Healthcare Providers:** Before purchasing ED medications, consult with a healthcare provider to get a legitimate prescription and guidance on where to buy genuine products. This reduces the risk of obtaining counterfeit medications and ensures proper treatment for your condition. **When Not to Use For Genuine Cialis:** **Heart Conditions:** Do not use Cialis if you have heart disease or are taking medications for chest pain (nitrates), as the combination can cause a dangerous drop in blood pressure. **Blood Pressure Issues:** Avoid Cialis if you have low or high blood pressure that is not controlled, as it can worsen these conditions. **Recent Stroke:** Do not use Cialis if you have recently had a stroke or a heart attack, as it may increase the risk of another event. **Eye Conditions:** Avoid Cialis if you have certain eye conditions, such as retinitis pigmentosa or a history of severe vision loss, as it can exacerbate these issues. **Allergies:** Do not use Cialis if you are allergic to Tadalafil or any other ingredients in the medication, as this can lead to serious allergic reactions. **For Counterfeit Cialis Black: General Avoidance:** Strongly advise against using counterfeit Cialis Black under any circumstances due to the unknown risks and potential for severe adverse effects. These pills can contain harmful or ineffective ingredients and may lead to serious health complications. Always opt for genuine medication obtained from reputable sources to ensure safety and effectiveness.

Warnings

For Genuine Cialis: **Interaction with Nitrates:** Do not take Cialis if you are using nitrates for chest pain, as this combination can cause a significant drop in blood pressure, which can be dangerous. **Health Conditions:** Cialis may not be suitable for everyone. People with heart problems, low blood pressure, or severe liver or kidney disease should consult their doctor before using it. **Medical Consultation:** Discuss any health concerns with your doctor before taking Cialis. Inform them about all the medications you are currently taking, including prescription drugs, over-the-counter medicines, and herbal supplements. **Alcohol Use:** Drinking alcohol can increase certain side effects of Cialis, such as dizziness and headache. It's best to limit alcohol consumption while using this medication. **For Counterfeit Cialis Black: High Risk of Adverse**

Effects: Using counterfeit pills carries a high risk of serious adverse effects. These pills may contain harmful substances, leading to severe health problems. Unknown Ingredients: The lack of regulation means that counterfeit Cialis Black can include unknown and potentially dangerous ingredients. Safety Concerns: The safety and effectiveness of counterfeit pills are not guaranteed. Always seek medications from legitimate and reputable sources to avoid these risks.

Dosage

For Genuine Cialis Black: Typical Dosage: Cialis Black is often marketed with a higher dosage, usually containing 80 mg of Tadalafil per pill. This is significantly higher than the regular Cialis doses. Administration: Take the pill with water, with or without food. Swallow the tablet whole without crushing or splitting it. Timing: It is recommended to take Cialis Black about 20 to 60 minutes before sexual activity. The effects can last up to 36 hours, providing a longer window of opportunity for sexual activity. Dosage Limitations: Due to the higher dosage, it is crucial not to take more than one pill in a 24-hour period to avoid the risk of severe side effects. Follow Doctor's Instructions: Always follow the dosage instructions provided by your healthcare professional. Do not exceed the recommended dose. Consult your doctor before adjusting the dosage to ensure it is safe for you. For Counterfeit Cialis Black: Avoid Usage: Do not attempt to determine or take any dosage of counterfeit Cialis Black. The lack of regulation means dosages can be incorrect or unsafe. Stick to genuine medication prescribed by a healthcare provider to ensure safety and effectiveness.

Interactions

For Genuine Cialis: Medications to Avoid: Nitrates: These are medications often prescribed for chest pain. Taking them with Cialis can cause a dangerous drop in blood pressure. Alpha-blockers: Used to treat high blood pressure or prostate conditions. Combining them with Cialis can lead to low blood pressure and dizziness. Antifungal Medications: Drugs like ketoconazole or itraconazole used to treat fungal infections. They can increase the levels of Cialis in your blood, leading to more side effects. Certain Antibiotics: Erythromycin, an antibiotic, can also increase Cialis levels in your blood and enhance side effects. Medications for High Blood Pressure: Taking these with Cialis can lower your blood pressure too much. HIV Protease Inhibitors: Used to treat HIV, these medications can increase Cialis levels in your blood and cause more side effects. Other ED Medications: Do not combine Cialis with other erectile dysfunction treatments to avoid excessive side effects. Inform Your Doctor: Always tell your healthcare provider about all the medications and supplements you are taking to avoid harmful interactions. Alcohol: Drinking alcohol while taking Cialis can increase side effects like dizziness and may reduce your ability to get an erection. For Counterfeit Cialis Black: High Risk of Dangerous Interactions: Counterfeit pills often have unknown ingredients, which can interact unpredictably with other medications. Unpredictable Effects: These interactions can cause severe health problems. Reputable Sources: Always get your medication from reputable sources to ensure it is safe and effective. Avoid counterfeit versions of Cialis Black.

FAQs

- **Is Generic Cialis Black safe for everyone?**

Generic Cialis Black is not suitable for everyone. Individuals with certain health conditions, particularly heart disease or those on medications containing nitrates, should avoid it. Consult a healthcare provider for personalized advice.

- **How long does the effect of Generic Cialis Black last?**

The effects of Generic Cialis Black can last up to 36 hours, which is longer than most other ED

medications. This extended duration offers a larger window of opportunity for sexual activity.

- **How should Generic Cialis Black be stored?**

Generic Cialis Black should be stored at room temperature, away from moisture and direct sunlight, and out of reach of children and pets.

- **Can I take Generic Cialis Black if I am taking other medications?**

It's important to inform your healthcare provider about all the medications you are currently taking as Generic Cialis Black can interact with certain drugs, particularly those used to treat heart conditions and high blood pressure.

- **Does Generic Cialis Black require a prescription?**

Yes, like other ED medications, Generic Cialis Black requires a prescription. Consult with a healthcare provider to determine if it is a suitable and safe option for you.

Other Details

For Genuine Cialis: With or Without Food: Cialis can be taken with or without food. Eating does not significantly affect how the medication works. **Alcohol Interaction:** While moderate alcohol consumption (such as a glass of wine) may be acceptable, excessive drinking can increase the risk of side effects like dizziness and can make it harder to achieve an erection. **Duration of Effects:** The effects of Cialis can last up to 36 hours, giving you a longer period during which you can achieve an erection when sexually aroused. **For Counterfeit Cialis Black: Avoid Use:** Due to the unknown risks associated with counterfeit pills, it is crucial to avoid their use entirely. These pills can contain harmful substances and incorrect dosages, posing serious health risks. **Consult a Doctor:** Always consult a healthcare provider to get a legitimate prescription and ensure you are using a safe and effective treatment for erectile dysfunction. **Verify Authenticity:** Make sure the medication is genuine by checking for official packaging, holograms, and approval seals. Purchase from reputable pharmacies to avoid counterfeit products. **Conclusion For Genuine Cialis: Summary:** Cialis is a proven and effective treatment for erectile dysfunction, helping men achieve and maintain an erection. It is important to use the medication as prescribed by a healthcare professional and be aware of potential interactions and side effects. **Consultation:** Always consult a doctor for personalized advice and to ensure safe usage of Cialis. Follow the prescribed dosage and avoid combining it with contraindicated medications. **For Counterfeit Cialis Black: Dangers:** Counterfeit Cialis Black poses significant health risks due to unknown ingredients and lack of regulation. These pills can cause severe side effects and potentially life-threatening reactions. **Seek Legitimate Treatment:** Always seek legitimate medical help for erectile dysfunction and obtain medications from reputable sources to ensure safety and effectiveness. **Responsible Use:** Advocate for safe and responsible use of medications by verifying their authenticity and consulting healthcare providers for proper guidance.

References

Mayo Foundation for Medical Education and Research. Tadalafil (Oral Route). In: Mayo Clinic. 2021. [Accessed 20 May 2024]. Available from: <https://www.mayoclinic.org/drugs-supplements/tadalafil-oral-route/description/drg-20067204> U.S. Food and Drug Administration. Medication Guide for Tadalafil. 2020. [Accessed 20 May 2024]. Available from: https://www.accessdata.fda.gov/drugsatfda_docs/label/2011/021368s20s21lbl.pdf WebMD LLC. Cialis (Tadalafil) Oral. In: WebMD. 2022. [Accessed 20 May 2024]. Available from: <https://www.webmd.com/drugs/2/drug-77881/cialis-oral/details> Drugs.com. Tadalafil. In: Drugs.com. 2023. [Accessed 20 May 2024]. Available from: <https://www.drugs.com/tadalafil.html>

Disclaimer

Disclaimer: The information on this site is provided for informational purposes only and is not medical advice. It does not replace professional medical consultation, diagnosis, or treatment. Do not self-medicate based on the information presented on this site. Always consult with a doctor or other qualified healthcare professional before making any decisions about your health.