

iMedix: Your Personal Health Advisor.

Gluco Burst 40 % Oral Gel Dextrose, Formulations

Search for medical details concerning Gluco Burst oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects This product usually has very few side effects. If you have any unusual effects from taking this product, tell your doctor or pharmacist promptly. If your doctor has directed you to use this product, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this product do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This product is used to treat low blood sugar levels. Symptoms of low blood sugar include sudden sweating, cold skin, shaking, irritability, fast heartbeat, hunger, blurred vision, dizziness, or tingling hands/feet. It is a good habit to carry glucose tablets, liquid, or gel to treat low blood sugar. If you don't have these reliable forms of glucose, rapidly raise your blood sugar by eating a quick source of sugar such as table sugar, honey, or candy, or drink fruit juice or non-diet soda. To help prevent low blood sugar, eat meals on a regular schedule, and do not skip meals. Check with your doctor or pharmacist about what to do if you miss a meal. **How to use Gluco Burst 40 % Oral Gel Dextrose, Formulations** Read and follow all directions on the product package before you need to take this product. Be sure to keep this product handy in case it is needed. If you have any questions, consult your doctor or pharmacist. Take this product by mouth as directed. If you are taking a chewable form of this product, chew it thoroughly before swallowing. Some liquid forms need to be shaken before use. Check the manufacturer's directions. Check your blood sugar level 10 to 15 minutes after taking this product. If your blood sugar level is still low, take another dose. The blood sugar level should be kept up by eating snacks such as crackers, cheese, a meat sandwich, or milk. If your symptoms do not get better within 20 minutes or if they get worse, or if you think you may have a serious medical problem, get medical help right away. **Precautions** Before taking glucose, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This product is safe to take during pregnancy when used as directed. It is unknown if this

product passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Does Gluco Burst 40 % Oral Gel Dextrose, Formulations interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Keep all regular medical and lab appointments. Attend a diabetes education program to learn more about how to manage your diabetes with medications, diet, exercise, and regular medical exams. Not applicable. Store at room temperature away from light and moisture. Do not store in the bathroom. Do not freeze liquid forms of this medication. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details
