

iMedix: Your Personal Health Advisor.

Haloperidol Lactate

Search for medical details concerning haloperidol lactate oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Dizziness, lightheadedness, drowsiness, difficulty urinating, sleep disturbances, headache, or anxiety may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Dizziness and lightheadedness can increase the risk of falling. Get up slowly when rising from a sitting or lying position. Tell your doctor right away if any of these side effects occur: muscle spasm/stiffness, shaking (tremor), restlessness, mask-like expression of the face, drooling/trouble swallowing, or shuffling walk. Your doctor may prescribe another medication to decrease these side effects. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication may rarely cause a condition known as tardive dyskinesia. In some cases, this condition may be permanent. Tell your doctor right away if you develop any facial/muscle twitching such as tongue thrusting, chewing movements, puffing or puckering of your mouth, or uncontrollable shaking. Tell your doctor right away if you have any serious side effects, including: nausea/vomiting that doesn't stop, stomach/abdominal pain, yellowing of eyes/skin, seizures, signs of infection (such as sore throat that doesn't go away, fever). Get medical help right away if you have any very serious side effects, including: slow heartbeat, severe dizziness, chest pain, fainting. This medication may rarely cause a very serious condition called neuroleptic malignant syndrome (NMS). Get medical help right away if you have any of the following symptoms: fever, muscle stiffness/pain/tenderness/weakness, severe tiredness, severe confusion, sweating, fast/irregular heartbeat, dark urine, signs of kidney problems (such as change in the amount of urine). In rare cases, haloperidol may increase your level of a certain chemical made by the body (prolactin). For females, this increase in prolactin may result in unwanted breast milk, missed/stopped periods, or difficulty becoming pregnant. For males, it may result in decreased sexual ability, inability to produce sperm, or enlarged breasts. If you develop any of these symptoms, tell your doctor right away. Rarely, males may have a painful or prolonged erection lasting 4 or more hours. If this occurs, stop using this drug and get medical help right away, or permanent problems could occur. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report

side effects to Health Canada at 1-866-234-2345. Uses Haloperidol is used to treat certain mental/mood disorders (such as schizophrenia, schizoaffective disorders). This medicine helps you to think more clearly, feel less nervous, and take part in everyday life. It can also help prevent suicide in people who are likely to harm themselves. It also reduces aggression and the desire to hurt others. It can decrease negative thoughts and hallucinations. Haloperidol can also be used to treat uncontrolled movements and outbursts of words/sounds related to Tourette's syndrome. Haloperidol is also used for severe behavior problems in hyperactive children when other treatments or medications have not worked. Haloperidol is a psychiatric medication (antipsychotic-type) that works by helping to restore the balance of certain natural substances in the brain (neurotransmitters).

How to use Haloperidol Lactate Take this medication by mouth with or without food as directed by your doctor. If you are using the liquid form of the medication, use the measuring dropper provided in the package to measure your dose. If you do not have the dropper, ask your pharmacist for a measured oral syringe so that your dose will be correct. The dosage is based on your medical condition and response to treatment. Use this medication regularly in order to get the most benefit from it. To help you remember, use it at the same time(s) each day. Do not stop taking this medication without consulting your doctor. Some conditions may become worse when the drug is suddenly stopped. Your dose may need to be gradually reduced. Tell your doctor if your symptoms do not improve or if they worsen.

Warnings There may be a slightly increased risk of serious, possibly fatal side effects (such as heart failure, fast/irregular heartbeat, pneumonia) when this medication is used by older adults with dementia. This medication is not approved for the treatment of dementia-related behavior problems. Discuss the risks and benefits of this medication, as well as other effective and possibly safer treatments for dementia-related behavior problems, with the doctor.

Precautions Before taking haloperidol, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: a certain severe nervous system problem (severe CNS depression), Parkinson's disease, bipolar disorder, difficulty urinating (for example, due to prostate problems), glaucoma, heart problems (such as angina), overactive thyroid (hyperthyroidism), seizures, low white blood cell count. Haloperidol may cause a condition that affects the heart rhythm (QT prolongation). QT prolongation can rarely cause serious (rarely fatal) fast/irregular heartbeat and other symptoms (such as severe dizziness, fainting) that need medical attention right away. The risk of QT prolongation may be increased if you have certain medical conditions or are taking other drugs that may cause QT prolongation. Before using haloperidol, tell your doctor or pharmacist of all the drugs you take and if you have any of the following conditions: certain heart problems (heart failure, slow heartbeat, QT prolongation in the EKG), family history of certain heart problems (QT prolongation in the EKG, sudden cardiac death). Low levels of potassium or magnesium in the blood may also increase your risk of QT prolongation. This risk may increase if you use certain drugs (such as diuretics/"water pills") or if you have conditions such as severe sweating, diarrhea, or vomiting. Talk to your doctor about using haloperidol safely. This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication may make you sweat less, making you more likely to get heat stroke. Avoid doing things that may cause you to overheat, such as hard work or exercise in hot weather, or using hot tubs. When the weather is hot, drink a lot of fluids and dress lightly. If you overheat, quickly look for a place to cool down and rest. Get medical help right away if you have a fever that does not go away, mental/mood changes, headache, or dizziness. Children may be more sensitive to the side effects of this drug, especially uncontrolled movements. Older adults may be more sensitive to the side effects of this drug, especially drowsiness, dizziness, lightheadedness, difficulty urinating, and heart effects such as QT prolongation (see above). Drowsiness, dizziness, and lightheadedness can increase the risk of falling. During pregnancy, this medication should be used only when clearly needed. Babies born to mothers who have used this drug during the last 3 months of pregnancy may rarely develop symptoms including muscle stiffness or shakiness, drowsiness, feeding/breathing difficulties, or constant crying. If you notice any of these symptoms in your newborn especially during their first month, tell the doctor right away. Since untreated mental/mood problems (such as schizophrenia, schizoaffective disorders) can be a serious condition, do not stop taking this medication unless directed by your doctor. If you are

planning pregnancy, become pregnant, or think you may be pregnant, immediately discuss with your doctor the benefits and risks of using this medication during pregnancy. This drug passes into breast milk and could have undesirable effects on a nursing infant. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: cabergoline, ketoconazole, lithium, methyldopa, drugs for Parkinson's disease (such as levodopa and carbidopa, selegiline), paroxetine, pergolide, quinupristin/dalfopristin, saquinavir. Many drugs besides haloperidol may affect the heart rhythm (QT prolongation), including amiodarone, dofetilide, pimozide, quinidine, sotalol, procainamide, macrolide antibiotics (such as amoxil, zithromax, flagyl, levofloxacin, ciprofloxacin), among others. Before using haloperidol, report all medications you are currently using to your doctor or pharmacist. Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely.

Side Effects

Uses

Interactions

Other Details
