

# iMedix: Your Personal Health Advisor.

## HEMORRHOIDAL Ointment With Applicator

---

Search for medical details concerning Hemorrhoidal (phenylephrine-mineral oil-petrolatum) rectal on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

**Side Effects** This medication does not usually have bothersome side effects when used as directed. Mild pain/stinging may occur if the rectal tissue is raw or bleeding. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Stop using this medication and tell your doctor right away if you have any serious side effects, including: fast/irregular heartbeat, pounding headache, nervousness, shakiness (tremor), trouble sleeping. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

**How to use HEMORRHOIDAL Ointment With Applicator** Follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. Use this product up to 4 times daily, usually in the morning and bedtime, or after each bowel movement or as directed by your doctor. Before use, clean the area with mild soap and water, rinse well, and pat dry. Remove the protective cover from the applicator tip, attach the applicator tip to the medication tube, and place a small amount of ointment on the applicator tip. Gently insert the applicator tip into the rectum and squeeze the tube to apply a small amount of this medication. Also apply some medication to the outside of the anus. After each use, clean the applicator tip well and replace the protective cover. The dosage is based on your medical condition and response to treatment. Do not use more often or for a longer time than recommended. If symptoms do not improve within 7 days, if bleeding/worsening pain occurs, or if you think you may have a serious medical problem, consult your doctor promptly.

**Precautions** Before using this product, tell your doctor or pharmacist if you are allergic to it; or to any of its ingredients; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. If you have any of the following health problems, consult your doctor or pharmacist before using this product: diabetes, heart problems (such as chest pain, heart attack), high blood pressure, overactive thyroid (hyperthyroidism), difficult urination due to blockage (for example, due to an enlarged prostate). Before having surgery, tell your doctor or dentist about all the

products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug are: MAO inhibitors (isocarboxazid, linezolid, metaxalone, methylene blue, moclobemide, phenelzine, procarbazine, rasagiline, safinamide, selegiline, tranylcypromine). Does HEMORRHOIDAL Ointment With Applicator interact with other drugs you are taking? Enter your medication into the iMedix interaction checker Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. To maintain normal bowel function and reduce the risk of hemorrhoids, eat a proper diet high in fiber, drink 6 to 8 glasses of water each day, and exercise regularly. If you are using this product on a regular schedule and miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Different brands of this medication may have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

---

## **Side Effects**

---

---

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

---

## Other Details

---

---